



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/325	A1	(11) International Publication Number: WO 99/51631
		(43) International Publication Date: 14 October 1999 (14.10.99)
(21) International Application Number: PCT/FI99/00290	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 April 1999 (06.04.99)		
(30) Priority Data: 980782 3 April 1998 (03.04.98) FI		
(71)(72) Applicants and Inventors: KORHONEN, Timo [FI/FI]; Huntutie 10 A, FIN-00950 Helsinki (FI). PALVA, Aini [FI/FI]; Vällitalontie 87 A, FIN-00660 Helsinki (FI). PALVA, Ilkka [FI/FI]; Vällitalontie 87 A, FIN-00660 Helsinki (FI). HYNÖNEN, Ulla [FI/FI]; Raisiontie 4 A 6, FIN-00280 Helsinki (FI). WESTERLUND-WIKSTRÖM, Benita [FI/FI]; Tähtelänkujä 5 A 3, FIN-10210 Inkoo (FI).	Published With international search report.	
(74) Agent: SEPPO LAINE OY; Itämerenkatu 3 B, FIN-00180 Helsinki (FI).		
(54) Title: A PROTEIN REGION RESPONSIBLE OF BINDING TO EPITHELIAL CELL TYPES AND A DNA SEQUENCE ENCODING SAID REGION		
(57) Abstract		
<p>This invention relates to a DNA molecule encoding a polypeptide responsible of binding to human and/or animal epithelial cell types. It has been found that various fragments of S-layer protein SlpA of <i>Lactobacillus brevis</i> has adhesive properties to epithelial cells types. It is possible to modify or improve the binding capacity of various prokaryotic or eucaryotic cells to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cell types by using lactobacillar surface structures of this invention. In particular, it is possible with the nucleotide sequences of this invention to improve the binding properties of a host cell having probiotic effects to human and/or animal epithelial cell types.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A PROTEIN REGION RESPONSIBLE OF BINDING TO EPITHELIAL CELL TYPES
AND A DNA SEQUENCE ENCODING SAID REGION

The present invention relates to DNA molecules encoding polypeptides responsible of
5 binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or
endothelial cells. Further this invention relates to vectors containing the DNA molecules and
hosts transformed with the DNA molecules of this invention.

The present invention relates also to a method of constructing new hosts or new proteins
10 capable of binding to human and/or animal epithelial cell types.

This invention also relates to genes encoding preselected proteins modified to bind to
human and/or animal epithelial cell types.

15 In addition this invention relates to a host cell, to a protein and to a method for carrying
preselected factors/properties to human and/or animal epithelial cells or cell surfaces by
using a *Lactobacillus brevis* strain or *L. brevis* SlpA protein.

Bacterial adhesion to human epithelial and subepithelial tissues is a decisive initial event in
20 successful colonization of tissue sites by invading bacteria. Several molecular ligand-
receptor interactions have been characterized for bacterial species that cause infectious
diseases in man or animals. Adhesion of pathogenic bacteria to the tissue at the infection
site helps the bacteria to resist mechanical defences of our body, such as peristalsis in the
intestine or flow of urine in the urinary tract. Adhesion is a key determinant in the host,
25 tissue and cell-type tropism of bacterial infections. Attachment to tissues is also important
for those bacteria that establish themselves as members of the normal bacterial flora in the
human body. Species of *Lactobacillus* are major members of the indigenous bacterial flora
in the gastrointestinal and the genital tract of man and animals. Lactobacilli are thought to
be beneficial to their host organism and have a long history of use in the gastrointestinal and

the urogenital tract to prevent or cure various minor illnesses. Probiotic effects of lactobacilli include exclusion of invasive pathogens from intestinal and vaginal surfaces, production of antimicrobial substances, stimulation of immune systems, as well as other physiological effects. As lactobacilli are members of the normal bacterial flora and food-grade organisms, their possible use as carriers of vaccine antigens in the intestine has aroused interest. The mechanisms by which lactobacilli bring about the probiotic effects have remained uncharacterized, but it is generally agreed that efficient adhesion to epithelial surfaces is important for the colonization of the intestine as well as for the effects associated with these bacteria. Isolates of lactobacilli have been shown to adhere to the intestinal epithelium of their hosts (Coconnier *et al.*, *Appl. Environ. Microbiol.* 58:2034 - 2039 (1992) and Henriksson *et al.*, *Appl. Environ. Microbiol.* 57:499 - 502 (1991)), but to date, however, no molecular ligand-receptor interactions of lactobacilli have been characterized. Considering the high number of lactobacilli in our body and the biotechnological, health-associated as well as ecological importance of these bacteria, molecular characterization of their adhesion mechanisms is important.

S-layers are paracrystalline surface protein arrays that are commonly expressed by species of Eubacteria and Archaeobacteria (reviewed in Messner and Sleytr, *Adv. Microb. Physiol.* 33: 213 -275 (1992) and Sleytr and Sára, *Trends Biotechnol.* 15:20-26 (1997). Most S-layers are composed of a single protein species, the S-layer protein, greatly varying in size in different bacterial genera. The S-layer subunits are very hydrophobic and crystallize to form a two-dimensional layer on the bacterial surface. The genes encoding S-layers are efficiently transcribed, and the S-layer protein is the dominant protein species representing 10-20% of the total cellular protein of the bacterial cell. Differing functions have been attributed to the S-layers of different bacterial species. These functions include maintenance of the cell shape, protection of cells from hostile environment, anchorage of extracellular enzymes to the bacterial cell wall and mediation of bacterial attachment to animal tissues (Chu *et al.*, *J. Biol. Chem.* 266 :15258-15265 (1991), Schneitz *et al.*, *J. Appl. Microbiol.* 74: 290- 294 (1993) and Toba *et al.*, *Appl. Environ. Microbiol.* 61:2467-2471 (1995)). The S-layer of the fish pathogen *Aeromonas salmonicida* binds to extracellular matrix proteins and increases bacterial virulence by promoting bacterial spread to cause systemic infection in the

fish (Chu *et al.*, *J. Biol. Chem.* 266:15258-15265 (1991)). Most S-layer proteins aggregate in physiological buffers and their functional analysis have been restricted to solid phase assays (Sleytr and Sára, *Trends Biotechnol.* 15:20-26 (1997)), which has remained a severe limitation in the functional analysis of these important surface proteins.

S-layers are expressed by various species of the genus *Lactobacillus* (Masuda and Kawata *FEMS Microbiol. Lett.* 20:145-150 (1983). Their role in bacterial adhesiveness to chicken epithelium as well as to human and mouse extracellular matrix have been suggested (Schneitz *et al.*, *J. Appl. Microbiol.* 74: 290- 294 (1993), Toba *et al.*, *Appl. Environ. Microbiol.* 61:2467-2471 (1995)), but overall, the functions of lactobacillar S-layers have remained poorly characterized. Primary structure of a few lactobacillar S-layers have been determined (Vidgrén *et al.* *J. Bacteriol.* 174: 7419 - 7427 (1992), Boot *et al.*, *J. Bacteriol.* 175: 6089- 6096 (1993) and Boot *et al.*, *J. Bacteriol.* 177: 7222- 7230. The predicted lactobacillar S-layer proteins are 40 to 50 kDa in molecular size and show similarity in amino acid compositions.

Lactobacilli are important bacterial colonizers of our intestinal surfaces. Despite their high number and potential symbiotic effects in our body, our knowledge of the colonization mechanisms that the lactobacilli use to attach and multiply our intestine have remained uncharacterized. This has been in part due to the restricted methodology to genetically manipulate these bacteria and also due to the lack of suitable methods to study the binding mechanisms of these bacteria to intestinal or other mucosal surfaces.

WO 97/14802 suggests the use of *Lactobacillus fermentum* 104R 29 kD adherence factor for promoting the activity of microorganism cells to bind to a receptor recognized on mucus. However, the finding of a factor capable of binding to mucus does not solve the problem of specifically carrying preselected factors to human or animal epithelial cells. Mucus on the surface of intestinal tract and any factors bound to the mucus are easily rinsed out from the intestine.

In this invention a DNA molecule encoding a protein region responsible of binding of the

protein to intestinal, urogenital, endothelial and/or other epithelial cell types has for the first time been identified and characterized. Although there has been some preliminary notions about the possible binding capability of the S-layer proteins of lactic acid bacteria (LAB) to human or animal cells, the binding property has not been confirmed to be due to the S-layer protein. The DNA molecule encoding a protein region responsible of binding was unexpectedly found from a gene encoding the S-layer protein of a species of lactic acid bacteria, *Lactobacillus brevis*. However, according to this invention a homologous DNA molecule encoding similar advantageous binding properties, may be synthetic or semisynthetic or originate from the same or another group of microorganisms.

This invention results in various advantages. This invention makes it for the first time possible to modify or improve the binding capacity of various prokaryotic or eukaryotic cells to human or animal epithelial cell types, like intestinal, urogenital and/or endothelial cell types by using lactobacillar surface structures. In particular, it is possible with the nucleotide sequences of this invention to improve the binding properties of a host cell having probiotic effects to human and/or animal epithelial cell types.

It is possible with the nucleotide sequences of this invention to modify or improve the binding properties of a preselected protein to human and/or animal epithelial cell types.

Furthermore it is possible by gene technological means to modify a host cell having or being modified to have the binding capability, to carry antigens, advantageously vaccine antigens to the intestinal and/or urogenital tract of humans and/or animals. In particular, it is possible to modify strains of *Lactobacillus brevis* species.

The probiotic properties of various strains of *Lactobacillus brevis* species can be improved by genetic modification. Totally new properties can be transferred to the strains of *Lactobacillus brevis* species or other hosts of this invention, which may carry these properties to human or animal gastrointestinal or urogenital tract.

Various host cells having or being modified to have the binding capability can be used to

colonize the human or animal gastrointestinal or urogenital tract and to exclude pathogens by binding to cell receptors, which would otherwise be bound by pathogens or by growing to cell surface areas which would otherwise be colonized by pathogens.

- 5 Next, the invention will be described in more detail with the aid of the attached figures, of which

Figure 1. Effect of S-layer removal on the adherence of *L. brevis* ATCC 8287 to human Intestine 407 cells. Panel a shows adherence of untreated bacteria to the intestinal cells, and
10 panel b shows the adhesiveness of bacteria treated with guanidine hydrochloride to remove S-layer from the bacterial surface.

Figure 2. Quantitative analysis of the effect of S-layer removal on the adherence of *L. brevis* to Intestine 407 cells. Panel A shows the number of adherent bacteria per epithelial cell; the
15 adhesion test with the treated and the untreated bacteria was performed at four different bacterial cell densities indicated below. Panel B shows the polyacrylamide gel electrophoresis in sodium dodecyl sulfate (SDS-PAGE) of the proteins released from the bacterial surface by the Laemmli sample buffer used in SDS-PAGE. The S-layer peptide is indicated by the arrow. The S-layer peptide is the dominant peptide species released from the cells but
20 not the only one.

Figure 3. Schematic representation of the SlpA fragments expressed as fusion to flagellin. On top, the hydropathicity plot of the entire 465-amino acid SlpA peptide, The most probable antigenic determinant is indicated by the dashed line, and the SlpA peptide has a
25 signal sequence of 30 amino acids. Below, the bars indicate the fragments expressed in *flhC*, the numbers refer to the N- and C-terminal amino acids in the SlpA peptide. Binding of the chimeric flagella to Intestine 407 (Int 407) cells is indicated on the right.

Figure 4. Binding of the SlpA96-200/ FlhC chimeric flagella to the human Intestine 407
30 (Panel a) and urinary bladder T24 cells (Panel c). The binding was visualized by indirect immunofluorescence. Panels b and d show the corresponding fields by light microscopy.

Panel e shows the binding of the Δ FliC flagella to Intestine 407 cells, and Panel g the control staining without flagella; the corresponding light microscopic fields are shown in Panels f and h. Arrows indicate binding of the chimeric flagella, arrowhead indicates lack of binding.

Figure 5. The nucleotide sequence of *Lactobacillus brevis* *SlpA* gene and the corresponding amino acid sequence. The first and last nucleotide residue of the coding sequence and of the various fragments is marked with an arrow. Correspondingly the first and last amino acid residue of the entire peptide and of the fragments is marked with a circle.

DNA molecules of the invention

In this invention a DNA molecule encoding a protein region responsible of binding to human and/or animal intestinal, urogenital, endothelial and/or other epithelial cell types has for the first time been identified and characterized.

In this invention it has been found that the S-layer protein SlpA of *Lactobacillus brevis* has adhesive properties to epithelial cell types. The primary structure of this protein and the corresponding gene has been described in WO 94/00581 and in Vidgren *et al.*, *J.*

Bacteriol. 174: 7419-7427 (1992). According to Palva, A in H. Bahl *et al.*, *FEMS Microbiology Reviews* 20 (1997):47 - 98 preliminary results have indicated that *L. brevis* S-layer protein could mediate binding to intestinal epithelial cells. However, until this date there has not been any suitable method to study the role of S-layer proteins of lactobacilli. For instance Schneitz *et al.* *J. Appl. Microbiol.* 74: 290- 294 (1993) have reported that the S-layer of *L. acidophilus* was involved in the adhesion of these bacteria to avian intestinal epithelial cells, but chemical removal of the S-layers did not affect the adhesion to Caco-2 cells. Guanidine hydrochloride extraction, which was used in the experiments, removes many protein types in addition to S-layer and hence the role of lactobacillar S-layers can not be deduced from these experiments.

By using a recently introduced flagella display system (Westerlund-Wikström *et al.* *Protein*

Engin. 10: 1319 - 1326 (1997)) we studied the adhesive properties of the S-layer protein SlpA of *Lactobacillus brevis*. We found, surprisingly, that the full length but also very short regions of the *Lactobacillus brevis* S-layer protein were capable of binding to epithelial cell types. The predicted size of the full-length S-layer protein is 465 amino acids. In the present invention various fragments of *slpA* were expressed as gene fusions in the variable region of the *fliC_H* gene of *Escherichia coli*. The resulting chimeric flagella were assessed by indirect immunofluorescence for binding to various epithelial cells.

The S-layer peptides needed for the binding were contained in fragments of 270, 215, 275, 150 and 105 amino acid residues respectively, the shortest fragment representing residues 96 through 200 in the S-layer protein. However, any fragment being a partial amino acid sequence of these sequences or of the entire *Lactobacillus brevis* S-layer protein and possessing similar binding capacity as the above mentioned fragments is a polypeptide of this invention and any DNA molecule encoding these polypeptides is a DNA molecule of this invention.

Chimeric flagella harbouring inserts that represented the N-terminal part of the S-layer protein bound to intestinal as well as urinary bladder cells, whereas the C-terminal part of the S-layer protein did not confer binding on the chimeric flagella. However, the C-terminal parts of the protein may have an effect in enhancing the efficiency of binding.

The S-layer expressing bacterium *Lactobacillus brevis* ATCC 8287 efficiently adhered to the human small intestinal cell line and to the human urinary bladder cell line. Bacterial adhesiveness to both cell lines was completely abolished after removal of the S-layer protein (SlpA) from the bacterial surface by guanidine hydrochloride extraction.

This invention is directed to a DNA molecule encoding a polypeptide capable of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cells. The cells may originate from human or animal origin, like from porcine or poultry origin or from pet animals. The cells may be normal or e.g. tumour cells. The DNA molecule may be a DNA molecule having the full length or the partial sequence i.e. the coding sequence

contained in the nucleotide sequence of any one of SEQ ID NO. 1, SEQ ID NO. 2, SEQ ID NO. 3, SEQ ID NO. 4, SEQ ID NO 5 or SEQ ID NO. 6, representing the various fragments of the *Lactobacillus brevis slpA* gene, excluding, however, the full length SEQ ID NO. 6.

5

By a partial nucleotide sequence is meant a nucleotide sequence lacking at least one nucleotide residue as compared to the nucleotide sequences of SEQ ID 1 to SEQ ID 6.

SEQ ID NO. 1 represents the 315 nucleotide residues from 286 to 600, SEQ ID NO. 2

- 10 represents the 450 nucleotide residues from 286 to 735, SEQ ID NO. 3 represents the 825 nucleotide residues from 286 to 1110, SEQ ID NO. 4 represents the 645 nucleotide residues from 91 to 735, SEQ ID NO 5 represents the 810 nucleotide residues from 91 to 900 and SEQ ID NO. 6 represents the entire coding sequence of *Lactobacillus brevis slpA* gene from 1 to 1395 nucleotide residues. The first and last nucleotide residue of the
15 nucleotide sequences of SEQ ID NO. 1 to SEQ ID NO. 6 is marked with an arrow in Figure 5.

The DNA molecule of this invention may be a DNA molecule encoding a polypeptide having the full length or the partial amino acid sequence i.e. the amino acid sequence

- 20 contained in any one of SEQ ID NO. 7, SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO.10, SEQ ID NO 11 or SEQ ID NO 12, representing the various fragments of *Lactobacillus brevis* SlpA protein, excluding, however, the full length SEQ ID NO. 12. SEQ ID NO. 7 represents the shortest amino acid sequence of 105 amino acid residues between 96 and 200, SEQ ID NO. 8 represents 150 amino acid residues between 96 and 245, SEQ ID NO.
25 9 represents 275 amino acid residues between 96 and 370, SEQ ID NO.10 represents 215 amino acid residues between 31 and 245 and SEQ ID NO 11 represents 270 amino acid residues between 31 and 300. SEQ ID NO 12 represents the amino acid sequence of the entire *Lactobacillus brevis* SlpA protein between 1 and 465. The first and last amino acid residue of the amino acid sequences of SEQ ID NO. 7 to SEQ ID NO. 12 is marked with a
30 circle in Figure 5.

By a partial amino acid sequence is meant an amino acid sequence lacking at least one amino acid compared to the amino acid sequences SEQ ID NO. 7 to SEQ ID NO. 12.

The present invention furthermore relates to DNA molecules, the sequences of which differ from the sequences of the above-identified molecules due to degeneracy of the genetic code, and which code for a polypeptide capable of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cells.

The present invention relates also to DNA molecules, the sequences of which hybridize to any one of the DNA molecules above encoding a polypeptide capable of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cells.

The term "hybridization" in this context means hybridization under conventional hybridization conditions, preferably under stringent conditions such as described by, e.g.

Sambrook *et al.* (1989, *Molecular Cloning, A Laboratory Manual* 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY). Typical stringent hybridization conditions are exemplified in example 7, but equal hybridizations can be carried out in slightly different conditions as is known to a person skilled in the art.

In example 7 the *slpA* gene of *L. brevis* has been hybridized to the chromosomal DNA of other *Lactobacillus* strains. As can be seen from example 7, the hybridization method is very useful and reliable method to find new DNA molecules of this invention. All the *L. brevis* strains tested gave positive hybridisation signal except two strains, which were shown to lack the S-layer protein. Other S-protein expressing lactobacilli belonging to other lactobacilli species gave negative result. *L. buchneri*, which is closely related to *L. brevis* gave also positive signal. *L. buchneri* carries S-layer and has the capability of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cells.

These nucleic acid molecules that hybridize to the nucleic acid molecules of the present

invention can in principle be derived from any organism possessing such nucleic acid molecules. Preferably, they are derived from lactic acid bacteria or bifidobacteria. Nucleic acid molecules hybridizing to the nucleic acid molecules of the present invention can be isolated, e.g., from genomic libraries or cDNA libraries of various organisms.

5

Such nucleic acid molecules can be identified and isolated by using the nucleic acid molecules of the present invention or fragments of these molecules or the reverse complements of these molecules, e.g. by hybridization according to standard techniques (see Sambrook *et al.*(1989)).

10

As hybridization probe, e.g. nucleic acid molecules can be used that have exactly or substantially the same nucleotide sequence indicated in the Figure 5 or fragments of said sequence. Preferably is used the entire nucleotide sequence of the coding sequence of the *slpA* gene. The fragments used as hybridization probes can also be synthetic fragments obtained by conventional synthesis techniques and the sequence of which is substantially identical to that of the nucleic acid molecules according to the invention. Once genes hybridizing to the nucleic acid molecules of the invention have been identified and isolated it is necessary to determine the sequence and to analyze the properties of the proteins coded for by said sequence.

20

The term "hybridizing DNA molecule" includes fragments, derivatives and allelic variants of the above-described nucleic acid molecules that code for the above-described protein (or its equivalent) or a biologically active fragment thereof. Fragments are understood to be parts of nucleic acid molecules long enough to code for the described protein (or its equivalent) or a biologically active fragment thereof. The term "derivative" means in this context that the nucleotide sequences of these molecules differ from the sequences of the above-described nucleic acid molecules in one or more positions and are highly homologous to said sequence.

25

30

The present invention is directed also to DNA molecules which are homologous with the DNA molecules contained in the coding sequences of any of SEQ ID 1 to SEQ ID 6 or with

the DNA molecules encoding a polypeptide having the amino acid sequence contained in any of SEQ ID 7 to SEQ ID 12 or with the degenerated forms of these DNA molecules.

"Homology" is understood to refer to a sequence identity of at least 50%, preferably more than 70% and still more preferably more than 90% on the length of at least 300 nucleotides. The deviations from the nucleic acid molecules described above can be the result of deletion, substitution, insertion, addition or combination.

Homology furthermore means that the respective nucleotide sequences or encoded proteins are functionally and/or structurally equivalent. The DNA molecules that are homologous to the DNA molecules described above and that are derivatives of said DNA molecules are regularly variations of said molecules which represent modifications having the same biological function. They may be naturally occurring variations, such as sequences of other organisms or mutations. These mutations may occur naturally or may be achieved by specific mutagenesis. Furthermore, these variations may be synthetically produced sequences.

The present invention furthermore relates to DNA molecules, the sequences of which have an amino acid sequence which shows at least 40 % identity, preferably at least 50 % identity, to a sequence contained above and which code for a polypeptide capable of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cells. The amino acid sequences of this invention show identity of less than 36% compared to any known amino acid sequence. When comparing the *L. brevis* slpA protein amino acid sequence with amino acid sequences of the prior art S-layer proteins known from lactobacilli, it was found that the highest identity was below 36 %.

By an amino acid sequence that is an "equivalent" of a specific amino acid sequence is meant an amino acid sequence that is not identical to the specific amino acid sequence, but rather contains at least some amino acid changes (deletion, substitutions, inversions, insertions, etc.) that do not essentially affect the biological activity of the protein as compared to a similar activity of the specific amino acid sequence, when used for a desired purpose. The biological activity of a polypeptide means here the capability of binding to epithelial cells.

Preferably, an "equivalent" amino acid sequence contains at least 40% - 99% identity at the amino acid level to the specific amino acid sequence, most preferably at least 50%, more preferably at least 60% and in an especially highly preferable embodiment, at least 95% identity, at the amino acid level.

5

The term "binding" is used here to mean the adherence of a cell, protein, protein region or polypeptide reasonably firmly to an epithelial cell, like intestinal, urogenital and/ or endothelial cell type. According to this invention the binding capacity has been measured by determining the binding of polypeptides encoded by DNA molecules expressed as gene
10 fusions in the variable region of the *fliC_{H7}* gene of *Escherichia coli*. The resulting chimeric flagella were assessed by indirect immunofluorescence for binding to human intestinal and human urinary bladder cells. The binding capacity was visually characterized to be very strong (++++), strong (+++), weak (+) or no binding at all (-). By binding is meant the adherence of proteins or cells to the epithelial cells, particularly to the surface of the cell.

15

The binding was specific: chimeric flagella harbouring inserts that represented the N-terminal part of the S-layer protein bound to intestinal as well as urinary bladder cell types, whereas the C-terminal part of the S-layer protein did not confer binding on the chimeric flagella. The S-layer expressing bacterium *Lactobacillus brevis* ATCC 8287 efficiently
20 adhered to the human small intestinal cell line and to the human urinary bladder cell line. Bacterial adhesiveness to both cell lines was completely abolished after removal of the S-layer protein (SlpA) from the bacterial surface by guanidine hydrochloride extraction.

In Table I is shown the differences in binding of different strains of lactic acid bacteria to
25 human epithelial cell line (Intestine 407).

Table I. Binding of different strains of lactic acid bacteria to human epithelial cell line Intestine 407.

Strains	Binding capacity
<i>L. acidophilus</i> JCM1132	-
<i>L. crispatus</i> JCM5810	+++*
A296-21	+
<i>L. amylovorus</i> F81	-
JCM5807	+
<i>L. gallinarum</i> T-50	-
<i>L. gasseri</i> JCM1130	+
<i>L. johnsonii</i> 5F49	+
<i>L. brevis</i> ATCC 8287	++++

* binding to the extracellular matrix secreted by the cell (see Toba *et al.*, *Appl. Environ. Microbiol.* 61:2467-2471 (1995))

As can be seen in Table I, from the lactobacilli tested only *L. brevis* binds strongly to epithelial cells, particularly to the surface of the cell.

By "lactic acid bacteria" are meant all Gram-positive; anaerobic, microaerophilic or aerotolerant; catalase negative; rods or cocci; most importantly they all produce lactic acid as sole, major or important product from the energy-yielding fermentation of sugars. In practice, genuine members of lactic acid bacteria include at least the following genera: *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Pediococcus*, *Carnobacterium*, *Sporolactobacillus*, *Streptococcus*, *Enterococcus*, *Aerococcus*, *Vagococcus*, *Tetragenococcus* and *Atopium*. Many characteristics typical to genuine LAB are also common to the genus *Bifidobacterium* which consists of important health-promoting intestinal bacteria.

Transfer of the binding property to new host cells

It is possible with the nucleotide sequences of this invention to modify or improve the binding capability of various prokaryotic or eukaryotic hosts to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cell types. The hosts cells are improved or modified to have the binding capability by transferring the host cells by at least one of the DNA molecules of this invention.

The binding capability may be transferred to any suitable bacterial host, for example to strains of lactic acid bacteria or bifidobacteria or to a fungal strain, like to a yeast strain.

A nucleotide sequence of this invention may be inserted into a DNA vector with conventional techniques, including blunt-ending or staggered-ending termini for ligation, restriction enzyme digestion to provide appropriate termini, filling in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulations are described in Sambrook *et al.* (1989, *Molecular Cloning, A Laboratory Manual* 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

To express a desired coding sequence, transcriptional and translational signals recognizable by the host are necessary. The nucleotide sequences of this invention may be operably linked to the transcriptional and secretory regulatory elements in an expression vector, and introduced into a host cell to produce desired protein under the control of such sequences. A DNA molecule is said to be capable of expressing a polypeptide if it contains expression control sequences which contain transcriptional regulatory information and such sequences are operably linked to the nucleotide sequence which encodes the desired polypeptide. An operable linkage is a linkage in which a sequence is connected to a regulatory sequence in such a way as to place expression of the sequence under the influence or control of the regulatory sequence.

Where the protein or protein region expression and secretion control sequences do not

function satisfactorily in a host cell, then sequences functional in the host cell may be substituted as necessary.

The vectors of the invention may comprise other operable linked functional elements such as

- 5 DNA elements which confer antibiotic resistance on a host cell, and which provide for an origin of replication, or for insertion of a desired sequence into the chromosome of a host cell.

- 10 To transform a host cell with the DNA constructs of the invention many vector systems are available depending upon whether it is desired to insert the desired protein's DNA construct into the host cell chromosomal DNA, or to allow it to exist in an extrachromosomal form.

- 15 Cells which have stably integrated the introduced DNA into their chromosomes are selected by also introducing one or more markers which allow selection of host cells which contain the expression vector in the chromosome, for example the marker may provide resistance to antibiotics. The selectable marker gene (that can be later removed by methods well known in the art) can either be directly provided on the same vector as that providing the desired DNA gene sequences to be expressed, or such markers may be introduced into the same cell by co-transformation.

- 20 Factors of importance in selecting a particular plasmid or phage vector include the ease with which recipient cells that contain the vector may be recognized and selected from those recipient cells, which do not contain the vector and the number of copies of the vector which are desired in a particular host.

- 25 After the introduction of the vector, recipient cells are grown in a selective medium, which selects for the growth of vector-containing cells. Expression of the nucleotide sequences of this invention result in the production of the desired protein, or the production of a fragment of this protein and/or a host having the desired properties.

- 30 The above mentioned DNA molecules including the entire gene of *L. brevis* *slpA* can be

expressed on the surface of other selected well known probiotic microbes in order to enhance and target their adhesion to the host cells to allow more efficient colonization.

Also the degree of the adhesion mediated by the polypeptides of this invention can be varied

- 5 by modifying their expression by different copy numbers, promoter strength regulation or by changing the length and amino acid composition of the binding domain with genetic engineering. With amino acid modifications encoded by the DNA molecules of this invention their host specificity may be affected if required.

- 10 Probiotics have been defined as live micro-organisms which beneficially affect the health of the host (human or animal) by improving its intestinal balance. To date, the term probiotic has been widened to include also live preparations used genitourinary to prevent infections and restore the disturbed microbiological ecological balance. Different characteristics associated with potentially health affecting bacteria may include e.g. i) acid and bile
15 stability, ii) adherence to intestinal cells, iii) colonization of the intestinal tract, iv) production of antimicrobial substances, v) antagonism against pathogenic bacteria and vi) safety in food and clinical use.

- According to this invention the probiotic effects of the hosts having (or modified to have)
20 the binding capacity to epithelial cells can be enhanced by genetic means.

- Various hosts of this invention can be used to colonize the human or animal gastrointestinal or urogenital tract and to exclude pathogens by binding to cell receptors, which would otherwise be bound by pathogens or by growing to cell surface areas, which would
25 otherwise be colonized by pathogens.

- As an example of establishing a surface expression of the polypeptides of this invention, one can apply well conserved cell wall binding anchors (e.g. C-terminal sequence of PrtP from *L. lactis* or other lactic acid bacteria, staphylococcal protein A, *Streptococcus*
30 pyogenes M6 protein, yeast anchor sequences etc.), fused to a spacer region and a DNA molecule of this invention which is preceded by a suitable promoter and signal sequence (see

e.g. Piard *et al. J. Bacteriol.* 179:3068-3072 (1997) and Steidler *et al. Appl. Environ. Microbiol.* 64: 342-345 (1998).

Enhancing properties of oral vaccine carriers

5

The above mentioned DNA molecules including the entire gene of *L. brevis slpA* can be transferred to another microorganism which could be for example a live vaccine carrier or a new putative host developed for vaccine purpose to enhance their adhesion properties.

- 10 As described above the degree of adhesion can be varied by modifying the expression of the DNA molecules of this invention by different copy numbers, promoter strength regulation or by changing the length and amino acid composition of the binding domain with genetic engineering. With amino acid modifications encoded by the DNA molecules of this invention their host specificity may be affected if required.

15

The portals of entry of many pathogens are mucosal surfaces and most infections also begin at some mucosal site. The natural immune response to a pathogen is also likely to begin at the mucosal site of entry. To induce such a response with a vaccine, a similar route for its delivery may be regarded as a rational approach. Most vaccines available to date are,

- 20 however, injected parenterally giving systemic response alone which may not be long lasting and protective enough. Oral vaccines are most likely to prevent intestinal diseases as they stimulate mucosal associated lymphoid tissue in the gastrointestinal tract directly. Thus, new strategies to prevent the adhesion, multiplication and invasion of pathogens at mucosal surfaces have become more acute and in some instances the only way to prevent
- 25 an infectious disease. Furthermore, the use of oral (or other mucosal) routes for immunization against infective diseases is desirable because oral vaccines are easier to administer, have higher compliance rates and are likely to be less expensive. Experimental data indicate that colonization is required for example for immune response when using recombinant *Streptococcus gordonii* expressing surface antigens (Fischetti *et al. Current*
- 30 *Opinion in biotechnology* 7:659-666 (1996) and Medagliani *et al. Proc.Natl.Acad.Sci USA* 92:6868-6872 (1995).

Mediating effective adherence to mucosal cells by the sequences of this invention, microorganisms carrying a homologous region to the identified binding region possess excellent properties as novel live oral vaccine carriers. Furthermore, it has been

- 5 documented that surface (S-) layers function as effective adjuvants. In addition to the binding function, the C-terminal part of the SlpA gene can be modified to carry desired antigen epitopes and to efficiently present them as multiple copies at the S-layer formed by the SlpA subunits. Either a heterogenous or uniform S-layer can be formed depending on whether the host carries both an antigen-expressing and wild-type *slpA* gene or only the
- 10 modified antigen-expressing gene. By this the amount of antigen on the cell surface may be affected. As described above, the vaccine antigen can also be expressed on the cell surface apart from a S-layer which in such a case functions as adhesion factor and adjuvant only.

Preferably expression hosts for oral vaccine carriers are strains of lactic acid bacteria and bifidobacteria, more preferably *L. brevis* species, in particular the strain *L. brevis* ATCC8287.

Transfer of the binding property to new proteins

- 20 In addition to the use of the polypeptides of this invention responsible of the binding property in living vaccine carriers, also non-living applications for these structures are fully possible. The desired antigen polypeptides and shorter antigen epitopes having been modified to have the binding property of this invention can be expressed in the host chosen and administered after isolation and purification as pure protein preparations. If required,
- 25 such preparations may be protected against degradation by using inert particles such as biodegradable microparticles, liposomes and cochelates (O'Hagan, *Novel Delivery Systems for Oral Vaccines*. CRC Press, Florida (1994)).

Applications of the molecules of this invention in bifunctional target-specific biomolecules

An unexplored but highly possible application for the DNA molecules of this invention are their use in bifunctional molecules i.e. by creating fusion constructs where the binding region mediates the adhesion to mucosal cells and the other domain functions as an active target-specific molecule.

Such bifunctionally acting molecules could be formed for example by combining the DNA molecules of this invention with DNA sequences encoding enzymes, single chain antibodies or pharmaceutical proteins or toxins.

Targeted enzymes could be applied for example in the gut for degradation of (i) lactose by β -galactosidase to decrease adverse effects in lactose intolerance or ii) milk proteins by proteases and peptidases e.g. to release bioactive peptides or to increase tolerance against milk allergy. Certain toxins bound specifically to the polypeptides of this invention could also be used for destruction of pathogens at different mucosal sites. The delivery of the bifunctional molecule could be by direct spray or liquid preparations (nasal mucosa), by melting capsules (vaginal mucosa) or by inert particles described above for vaccine delivery systems (gastrointestinal tract).

With single chain antibodies and pharmaceutical proteins linked to the binding regions of this invention unlimited amounts of applications for mucosal prevention and medication of infectious diseases can be predicted.

Use of *Lactobacillus brevis* as a probiotic host

According to this invention *L. brevis* as such may function as a novel probiotic strain particularly due to its highly efficient binding capacity both in the gut and urinary tract even though its use in practical applications is yet unraveled. Its probiotic effect can be further enhanced for example by introducing genes encoding i) production of selected

bacteriocin(s) and the corresponding immunity and ii) other antimicrobial substances to antagonist pathogens and iii) enzymes increasing its metabolic activity towards available substrates to strengthen its competitiveness in the chosen niche.

5 Also the degree of the SlpA mediated adhesion can be varied by modifying its expression by different copy numbers, by promoter strength regulation or by changing the length and amino acid composition of the binding domain with genetic engineering. With amino acid modifications of the SlpA binding domain their host specificity may be affected if required.

10 In addition to the demonstrated *slpA* encoded binding capacity of *L. brevis* ATCC8287, this strain is also highly resistant in low pH and bile thus possessing the key characteristics of a probiotic strain. *L. brevis* strains have also been shown to have antagonistic effects against intestinal pathogens. Furthermore, *L. brevis* can be regularly found in the intestine of man and animals as well as in a variety of fermented food products.

15

The following examples and figures provide further details of the invention.

Example 1

20 *Bacteria*-The strain ATCC 8287 of *L. brevis* and its *slpA* gene have been described (Vidgrén *et al. J. Bacteriol.* 9:7419-7427 (1992)). The bacteria were grown overnight at 37 °C in 20 ml of static MRS broth. To extract the S-layer (Masuda, K. & Kawata, T. *Microbiol. Immunol.* 23:941-953 (1979)), the bacterial cells were washed once with phosphate buffered saline, pH 7.1 (PBS) and once with distilled water, and the cells were
25 then suspended in 0.8 ml of distilled water. The cells were then incubated for 2 h at 37 °C with 7.2 ml of 2 M guanidine hydrochloride; the control cells were incubated similarly in PBS alone. The guanidine hydrochloride-treated cells were washed once with 2 M guanidine hydrochloride and then twice with PBS; the control cells were washed twice with PBS. For adherence assays, the bacterial cells were suspended in the cell culture medium used with
30 the target epithelial cell lines (see below).

Bacterial adherence assays—The bacterial adhesion to cultured human epithelial cells was evaluated essentially as described earlier (Tarkkanen *et al.*, *Infect. Immunol.* 65:1546-1549 (1997)). The human small intestine Intestine 407 (ATCC CCL6) cells were cultivated to confluence in RPMI 1640 medium (Gibco Laboratories, Grand Island, N.Y.) supplemented with 10 % (w/v) fetal calf serum (PAA Laboratories GmbH, Linz, Austria), 1 % (w/v) L-glutamine (Life Technologies), 1 % (w/v) nonessential amino acids (Gibco Laboratories, Grand Island, N.Y.), and gentamycin (50 µg/ml). The human urinary bladder transitional T24 cells (ATCC HTB-4) were cultured in McCoy's 5A medium (Life Technologies) supplemented with fetal calf serum, L-glutamine, and gentamycin as above. The cell lines were cultured on diagnostic glass slides (Knittel Glassbearbeitungs GmbH, Braunschweig, Germany). Before the adhesion assays, the cells were washed once with PBS. The bacteria were suspended in RPMI 1640 medium at the concentrations ranging from 5×10^7 to 10^9 cells/ml, and 40 µl of the suspension per well was added to the epithelial cells and the slides were incubated for 1 h at 37 °C in a moist chamber. The slides were washed five times at room temperature with PBS for five min each and fixed for 10 min with methanol. The cells with adherent bacteria were then examined in a BX50 microscope (Olympus Optical Co., Hamburg, Germany) either directly by Nomarski interference optics (for photographing) or, for quantitative analysis, stained for 5 min with 10 % v/v Giemsa stain and analyzed by light microscopy. The number and standard deviation of adherent bacteria on 20 epithelial cells was calculated. In inhibition studies, purified Fab fragments from anti-SlpA immunoglobulins (Vidgrén *et al. J. Bacteriol.* 9:7419-7427 (1992)) or from control immunoglobulins raised against fimbrial proteins of *Escherichia coli* (Westerlund *et al. Zbl. Bakt.* 278:229-237 (1993)) were incubated with the bacterial suspensions for 30 min over crushed ice prior to the adhesion assay, the control bacteria were incubated similarly but in PBS alone. The Fab fragments were prepared by a routine procedure (Porter, *Biochem. J.* 73:119-126 (1959)) and tested at the final concentration of 500 µg/ml.

Example 2

Flagella display—The principle of the flagella display system used here was recently described (Westerlund-Wikström *et al.*, *Prot. Engin.* 10:1319-1326 (1997)). Fragments

representing different parts of the *slpA* gene were amplified by polymerase chain reaction (PCR) with *Pfu* polymerase and using chromosomal DNA from the *L. brevis* strain ATCC 8287 as the template. The primers were designed on the basis of the nucleotide sequence of *slpA* (Vidgrén *et al. J. Bacteriol.* 9:7419-7427 (1992)) and contained an *AccI* restriction site at the 5' termini. The primers encoded the SlpA peptide sequences 31-245 (SEQ ID NO. 10; this peptide sequence is encoded by the nucleotide sequence SEQ ID NO. 4, which represents the nucleotide residues from 91-735), 31-300 (SEQ ID NO. 11 this peptide sequence is encoded by the nucleotide sequence SEQ ID NO. 5, which represents the nucleotide residues from 91-900), 96-200 (SEQ ID NO. 7; this peptide sequence is encoded by the nucleotide sequence SEQ ID NO. 1, which represents the nucleotide residues from 286-600), 96-245 (SEQ ID NO. 8; this peptide sequence is encoded by the nucleotide sequence SEQ ID NO. 2, which represents the nucleotide residues from 286-735), 96-370 (SEQ ID NO. 9; this peptide sequence is encoded by the nucleotide sequence SEQ ID NO. 3, which represents the nucleotide residues from 286-1110), or 239-447 (SEQ ID NO. 13; this peptide sequence is encoded by the nucleotide sequence SEQ ID NO. 14, which represents the nucleotide residues from 715-1341), where the residue numbers include the 30-mer signal sequence of the SlpA peptide. The *slpA* fragments were cloned into the *AccI* site in the plasmid pFliC_{77Δ} deleted for 174 bp in the variable region of *fliC* and expressed in trans in *E. coli* that is *fliC::Tn10* and *fimA::cat* (Westerlund-Wikström *et al. Prot. Eng.* 10:1319-1326(1997)). The flagellar filaments were extracted and, after a sodium dodecylsulfate gel electrophoresis (SDS-PAGE), adjusted to an equal concentration of the FliC peptide as described recently (Westerlund-Wikström *et al., Prot. Engin.* 10:1319-1326 (1997)). The flagella lacking an insert, i.e. the ΔFliC filaments, were available from previous work (Westerlund-Wikström *et al.* (1997)).

Example 3

Binding tests with chimeric flagella-The binding of the chimeric flagella onto the epithelial cells was assessed by indirect immunofluorescence as detailed recently (Westerlund-Wikström *et al.* (1997)). Briefly, the epithelial cells were washed at room temperature with PBS, fixed with methanol for 10 min at -20 °C, and then washed with PBS at room

temperature. The flagellar extracts (40 μ l; 20 μ g/ml in PBS) were added and the slides were kept for 5 h at 4°C. After washing and a second fixing with methanol, the bound flagella were visualized by staining with immunoglobulin G molecules from an anti-H7-flagella rabbit antiserum and with fluorescein isothiocyanate-labelled secondary antibodies as detailed (Westerlund-Wikström *et al.* (1997)). The control assays included staining of the epithelial cells as above but using the Δ FliC flagella lacking an insert, or omitting the flagellar extract, or the flagellar extract and the immunoglobulins in the staining procedure.

Immunological methods-For immuno electron microscopy, bacterial cells expressing the various flagellar constructs were suspended in Luria broth and immobilized on copper grids coated with Pioloform and carbon. The bacteria were left to react with an anti-H7-flagella antiserum (Westerlund-Wikström *et al.* (1997)); diluted 1/300 in PBS containing 10 mg/ml BSA or with an anti-SlpA antiserum (Vidgrén *et al. J. Bacteriol.* 9:7419-7427 (1992)); diluted 1/300 in PBS-BSA) for 90 min at room temperature. The grids were washed in PBS containing 1 mg/ml BSA, and the bound antibodies were detected with Auoprobe™EM Protein A-conjugate (Amersham, Amersham Place; Little Chalfont, Buckinghamshire, UK; diluted 1/40). The grids were examined in a Jeol JEM-100CX transmission electron microscope at an operating voltage of 60 kV. For immunoblotting, flagellar preparations were analyzed by SDS-PAGE using a 1 % (w/v) stacking gel and a 10 % separating gel. Polypeptides were transferred onto a nitrocellulose membrane using a semi-dry transfer apparatus at 0.9 mA/cm² membrane for 2 h at 4 °C. After transfer, the membrane was quenched with PBS containing 20 mg/ml BSA for 16 h at 20 °C and washed with PBS. Polypeptides were visualized by staining with diluted polyclonal anti-flagella antibodies or anti-SlpA antibodies and alkaline-fosfatase-conjugated secondary antibodies as described (Westerlund-Wikström *et al.* (1997)). A phosphatase substrate solution containing nitroblue-tetrazolium (162 μ g/ml) and 5-bromo-4-chloro-3-indolyl-1-phosphate (370 μ g/ml) was used.

Example 4

Adherence of L. brevis ATCC 8287 to intestinal cells-We initially assessed the adhesiveness

of the *L. brevis* strain ATCC 8287 to the human small intestine cell line Intestine 407. The strain showed an efficient adhesion to the intestinal cells (Fig. 1A). The cells of *L. brevis* ATCC 8287 express the S-layer protein SlpA as their major cell surface protein (Vidgrén *et al. J. Bacteriol.* 9:7419-7427 (1992)), and we therefore extracted the S-layer from the bacterial surface and determined how this affected the adhesion. Extraction of cells with 2M guanidine hydrochloride is a routine procedure to remove bacterial S-layers, and the treatment does not lyse the bacterial cells. Treatment with guanidine hydrochloride completely abolished the adhesiveness of *L. brevis* (Fig. 1B). The S-layer peptide was the dominant peptide species released from the cells (Fig. 2B), and the results suggested a role for SlpA in bacterial adhesiveness.

Example 5

Expression of slpA fragments in fliC-The *slpA* gene encoding the S-layer protein of *L. brevis* ATCC 8287 has been described (Vidgrén *et al. J. Bacteriol.* 9:7419-7427 (1992)). We cloned fragments of *slpA* into the *AccI* site in the plasmid pFliC_{HTA} that contains a 174-bp deletion in the variable region of *fliC*, and the chimeric flagella were expressed in *E. coli* JT1 that is *fliC::Tn10* and *fimA::cat*. Schematic presentation of the *slpA* fragments expressed by the flagella display are shown in Fig.3. Western blots of the flagellar preparations with anti-H7 and anti-SlpA polyclonal antibodies showed that the apparent size of the chimeric flagellins corresponded to those predicted from the nucleotide sequence, i.e. it increased with the size of the insert in *AfliC* (data not shown). The polypeptides of smaller size that were present in the preparations and also reacted with the antibodies most likely were flagellar minor proteins (as noted earlier; Westerlund-Wikström *et al., Prot. Engin.* 10:1319-1326 (1997)) or degradation products. We noted by electron microscopy (data not shown) that the chimeric flagella expressing the larger inserts (> 200 amino acids) had short flagellar filaments and thus these preparations were reduced in the relative amount of FliC as compared to the hook and cap proteins of the flagella. The chimeric flagella SlpA31-245/ΔFliC, SlpA31-300/ΔFliC, SlpA96-370/ΔFliC and SlpA239-447/ΔFliC reacted with both the anti-H7 and the anti-SlpA antibodies, whereas the flagella with the two shortest inserts in the constructs SlpA96-245/ΔFliC and SlpA96-200/ΔFliC reacted with the anti-H7

antibodies and showed the expected apparent molecular size but failed to react with the anti-SlpA antiserum in Western blots.

Example 6

5

Binding of chimeric flagella to epithelial cells-We next analyzed by an indirect immunofluorescence assay the binding of the chimeric flagella to Intestine 407 cells, representative examples of the assays are shown in Fig. 4. The SlpA31-300/ Δ FliC construct, carrying the N-terminal part of the molecule, exhibited binding to the Intestine 407 cells, whereas the construct SlpA239-447/ Δ FliC representing the C-terminal part, failed to bind (data not shown). No binding was observed with the Δ FliC construct lacking an insert. These results indicated that the binding region is located within the N-terminal part of the SlpA molecule. We constructed and tested various fragments covering different regions of the N-terminus, and the shortest fragment supporting adhesion to Intestine 407 cells was the construct SlpA96-200/ Δ FliC that contained an 105 amino acid-long insert (Fig. 4A).

15

We also tested the binding of the SlpA96-200/ Δ FliC construct to the human urinary bladder cell line T24 (Fig. 4C). The chimeric flagella bound efficiently to the urinary bladder cells, whereas no binding of the Δ FliC flagella were observed. The *L. brevis* ATCC 8287 cells exhibited an efficient adhesion to the urinary bladder cell line, and the adhesiveness was abolished after treatment of the bacteria with guanidine hydrochloride (data not shown). It was also shown that *L. brevis* ATCC 8277 cells bound to CaCo-2 cells.

20

Example 7

25

Hybridization of the slpA gene of L. brevis ATCC8287 to other Lactobacillus strains-

The aim of this experiment was to study the presence of *L. brevis* slp-gene homologs in other *Lactobacillus* strains. For this purpose chromosomal DNA was isolated from the following *L. brevis* strains: all the six strains available in the DSM culture collections (DSM 20054, 1267, 1268, 1269, 2647, 2647, 6235), five strains of the Japanese collection (Yasui *et al.* *FEMS Microbiology Letter* 133:183-186 (1995), one strain (VK3) from the culture

30

collection of TNO (delivered by prof. Peter Powels), one strain of the University of Cornell, U.S.A. (delivered by prof. Carl Batt) and one own isolate from pig intestine identified by API as *L. brevis*/*L. buchneri*. In addition, several chromosomal DNA samples of S-layer carrying *Lactobacilli* isolated from pig intestine and identified by API as putative members of *L. acidophilus*, *L. fermentum*, *L. crispatus* and *L. delbrueckii* species were tested.

Hybridization was performed in the following buffer: 5xSSC, 1% Blocking Reagent (Boehringer Mannheim), 0,02% SDS, 0,1 % lauryl sarcosine at 68°C overnight using the full length *L. brevis* *slpA* gene labelled with DIG (digoxigenin- dUTP, Boehringer Mannheim) as probe. Hybridization was followed by washing steps in decreasing salt concentration and increasing temperature. The final wash step was in 0.1xSSC, 0.1% SDS at 68°C for 15 min twice. The negative control used was an equal amount of *E.coli* or calf thymus DNA to that of test DNA.

All *L. brevis* strains tested gave a positive hybridization signal except the strain VK3 from TNO and the Japanese strain Yasui 0296961015. These two hybridization negative *L. brevis* strains were also shown to lack the S-layer protein by SDS-PAGE analysis. Furthermore, the other above mentioned S-layer expressing lactobacilli not belonging to the *L. brevis* and/or *L. buchneri* group remained negative in the hybridization test.

We have also shown previously in WO 94/00581 that the chromosomal DNA of *L. buchneri* strain DSM 20057 strongly hybridizes with the *slp*-gene probe used above under stringent conditions, whereas S-layer protein carrying *L. helveticus* and *L. acidophilus* strains remains hybridization negative under same conditions.

We have also shown that the *L. brevis*/*L. buchneri* strains hybridizing with the ATCC8287 *slp*-probe also effectively bind to the Intestine 407 cells whereas non-S-layer carrying *L. brevis* strains do not adhere to Intestine 407 cells.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Korhonen et al.
(B) STREET: Huntutie 10 A
(C) CITY: Helsinki
(E) COUNTRY: Finland
(F) POSTAL CODE (ZIP): 00950

(ii) TITLE OF INVENTION: A protein region responsible of binding to epithelial cell types and a DNA sequence encoding said region

(iii) NUMBER OF SEQUENCES: 16

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: FI 980782
(B) FILING DATE: 03-APR-1998

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 315 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GTTAAGACCA CTAACCGTGG TTCAGTTTAC TACCGTGTGG TAACGATGGA TGGCAAGTAC	60
CGTGGTTACG TTTATGGTGG CAAGTCTGAC ACTGCCTTTG CTGGTGGTAT CAAGTCTGCT	120
GAAACGACTA CTAAGGCTGA TATGCCTGCA CGTACTACTG GGTCTACTTT AACTGACACT	180
TCAAAGAACA CTCTTTGAC GGCTCCTAAG TAACTCAAT ACAAGGCAAG TAAAGTTAGC	240
CTTTATGGTG TTGCTAAGGA CACCAAGTTT ACTGTAGATC AGGCTGCTAC TAAGACTCGT	300
GAAGGTTTAT TATAC	315

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 450 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

5 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

10	GTTAAGACCA CTAACCGTGG TTCAGTTTAC TACCGTGTG TAACGATGGA TGGCAAGTAC	60
	CGTGGTTACG TTTATGGTGG CAACTCTGAC ACTGCCTTTG CTGGTGGTAT CAAGCTCGCT	120
	GAAACGACTA CTAAGGCTGA TATGCCTGCA CGTACTACTG GGTCTACTT AACTGACACT	180
15	TCAAAGAACA CTCITTTGAC GGCTCCTAAG TACTACTCAAT ACAAGGCAAG TAAAGTTAGC	240
	CTTTATGGTG TTGCTAAGGA CACCAAGTTT ACTGTAGATC AGGCTGCTAC TAAGACTCGT	300
20	GAAGGTTTAT TATACTATCA CGTAACTGCT ACTAACGGTA GTGGTATTAG TGGTTGGATT	360
	TACGCTGGTA AGGGCTTCAG TACTACTGCT ACTGGTACAC AAGTACTTGG TGGTCTGTCA	420
	ACTGATAAGT CAGTTACAGC AACCAACGAT	450

25 (2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 825 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: DNA (genomic)

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

	GTTAAGACCA CTAACCGTGG TTCAGTTTAC TACCGTGTG TAACGATGGA TGGCAAGTAC	60
40	CGTGGTTACG TTTATGGTGG CAACTCTGAC ACTGCCTTTG CTGGTGGTAT CAAGCTCGCT	120
	GAAACGACTA CTAAGGCTGA TATGCCTGCA CGTACTACTG GGTCTACTT AACTGACACT	180
	TCAAAGAACA CTCITTTGAC GGCTCCTAAG TACTACTCAAT ACAAGGCAAG TAAAGTTAGC	240
45	CTTTATGGTG TTGCTAAGGA CACCAAGTTT ACTGTAGATC AGGCTGCTAC TAAGACTCGT	300
	GAAGGTTTAT TATACTATCA CGTAACTGCT ACTAACGGTA GTGGTATTAG TGGTTGGATT	360
	TACGCTGGTA AGGGCTTCAG TACTACTGCT ACTGGTACAC AAGTACTTGG TGGTCTGTCA	420
50	ACTGATAAGT CAGTTACAGC AACCAACGAT AACAGTGTTA AGATTGTTTA CCGTACGACT	480
	GATGGCACTC AAGTTGGTTC TAACACTTGG GTAACCTCAA CTGATGGTAC AAAGGCAAGT	540
55	TCTAAGGTAA GCGATAAGGC CGCCGATCAA ACTGCTCTTG AAGCCTACAT CAATGCTAAC	600
	AAGCCTAGCG GTTACTACTG AACTAACCTT AATGCTGCAG ATGCTACCTA TGGTAACACA	660

29

GTTTACGCTA CTGTTTCCCA AGCAGCTACT TCTAAGGTCG CTTTGAAGGT CTCAGGGACT 720
 CCTGTTACTA CTGCATTGAC TACAGCTGAT GCTAATGATA AGGTTGCAGC TAAAGATACC 780
 5 ACTGCTAATG GTAGTTCCTGT TGCAGGCTCA ACAGTCTATG CTGCT 825

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 645 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

20 AAGTCATACG CTACTGCAGG TGCCTATTCA ACGTTAAAGA CGGACGCTGC TACTCGTAAC 60
 GTCGAAGCTA CTGGTACTAA CGCTTTATAC ACGAAGCCAG GTACTGTTAA GGGTGCTAAG 120
 GTTGTGCTT CTAAGGCTAC TATGGCTAAG TTAGCTTCTT CAAAGAAGTC AGCTGACTAC 180
 25 TTCCGTGCTT ACGGTGTTAA GACCACTAAC CGTGGTTCAG TTTACTACCG TGTTGTAAAC 240
 ATGGATGGCA AGTACCGTGG TTACGTTTAT GGTGGCAAGT CTGACACTGC CTTTGTCTGGT 300
 30 GGTATCAAGT CTGCTGAAAC GACTACTAAG GCTGATATGC CTGCACGTAC TACTGGGTTT 360
 TACTTAACTG ACACTTCAAA GAACACTCTT TGGACGGCTC CTAAGTACAC TCAATACAAG 420
 GCAAGTAAAG TTAGCCTTTA TGGTGTGCT AAGGACCA AGTTTACTGT AGATCAGGCT 480
 35 GCTACTAAGA CTCGTGAAGG TTCATTATAC TATCACGTAA CTGCTACTAA CGGTAGTGGT 540
 ATTAGTGGTT GGATTTCAGC TGTAAAGGCG TTCAGTACTA CTGCTACTGG TACACAAGTA 600
 40 CTTGGTGGTC TGCTAACTGA TAAGTCAGTT ACAGCAACCA ACGAT 645

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 810 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

50 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

55 AAGTCATACG CTACTGCAGG TGCCTATTCA ACGTTAAAGA CGGACGCTGC TACTCGTAAC 60
 GTCGAAGCTA CTGGTACTAA CGCTTTATAC ACGAAGCCAG GTACTGTTAA GGGTGCTAAG 120

GTTGTCGCTT CTAAGGCTAC TATGGCTAAG TTAGCTTCCT CAAAGAAGTC AGCTGACTAC 180
 TTCCGTGCTT ACGGTGTTAA GACCACTAAC CGTGGTTCAG TTACTACCG TGTTGTAACG 240
 15 ATGGATGGCA AGTACCGTGG TTACGTTTAT GGTGGCAAGT CTGACACTGC CTTTGCTGGT 300
 GGTATCAAGT CTGCTGAAAC GACTACTAAG GCTGATATGC CTGCACGTAC TACTGGGTTTC 360
 TACTTAACTG ACACCTTCAA GAACACTCTT TGGACGGCTC CTAAGTACAC TCAATACAAG 420
 10 GCAAGTAAAG TTAGCCTTTA TGGTGTGCT AAGGACACCA AGTTTACTGT AGATCAGGCT 480
 GCTACTAAGA CTCGTGAAGG TTCATTATAC TATCAGTAA CTGCTACTAA CGGTAGTGST 540
 15 ATTAGTGGTT GGATTTACGC TGGTAAGGGC TTCAGTACTA CTGCTACTGG TACACAAGTA 600
 CTGGTGGTC TGTCACCTGA TAAGTCAGTT ACAGCAACCA ACGATAACAG TGTTAAGATT 660
 GTTTACCGTA CGACTGATGG CACTCAAGTT GGTTCCTAACA CTGGGGTAAC TTCAACTGAT 720
 20 GGTACAAAGG CAGGTCTCTAA GGTAAGCGAT AAGGCCGCCG ATCAAAGTGC TCTTGAAGCC 780
 TACATCAATG CTAACAAGCC TAGCGGTTAC 810

25 (2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1395 base pairs
 (B) TYPE: nucleic acid
 30 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: DNA (genomic)

35

(x) PUBLICATION INFORMATION:

- (A) AUTHORS: Vidgren, G
 Palva, I
 Pakkanen, R
 40 Lounatmaa, K
 Palva, A
 (B) TITLE: S-Layer Protein Gene of *Lactobacillus brevis*:
 Cloning by Polymerase Chain Reaction and
 Determination of the Nucleotide Sequence
 45 (C) JOURNAL: J. Bacteriol.
 (D) VOLUME: 174
 (E) ISSUE: 22
 (F) PAGES: 7419-7427
 (G) DATE: 1992
 50 (K) RELEVANT RESIDUES IN SEQ ID NO: 6: FROM 1 TO 1395

(x) PUBLICATION INFORMATION:

- (H) DOCUMENT NUMBER: WO 94/00581 A1
 (I) FILING DATE: 24-JUN-1993
 55 (J) PUBLICATION DATE: 06-JAN-1994

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

	ATGCAATCAA GTTTAAAGAA ATCTCTTAC TTGGCCCTTG CCGCATTGAG CTTTGCTGGT	60
	GTGTGCTCCG TTCAACGAC TGCTTCAGCT AAGTCATACG CTACTGCAGG TGCCTATTCA	120
5	ACGTTAAAGA CGGACGCTGC TACTCGTAAC GTCGAAGCTA CTGGTACTAA CGCTTTATAC	180
	ACGAAGCCAG GTACTGTTAA GGGTGCTAAG GTTGTGCTT CTAAGGCTAC TATGGCTAAG	240
10	TTAGTCTCTT CAAAGAAGTC AGCTGACTAC TTCCGTGCTT ACGGTGTATA GACCACTAAC	300
	CGTGGTTCAG TTACTACCG TGTGTAAAG ATGGATGGCA AGTACCGTGG TTACGTTTAT	360
	GGTGGCAAGT CTGACACTGC CTTTGCTGGT GGTATCAAGT CTGCTGAAAC GACTACTAAG	420
15	GCTGATATGC CTGACGCTAC TACTGGGTTT TACTTAACTG ACACCTTCAA GAACACTCTT	480
	TGGACGGCTC CTAAGTACAC TCAATACAAG GCAAGTAAAG TTAGCCTTTA TGGTGTGTCT	540
20	AAGGACACCA AGTTTACTGT AGATCAGGCT GCTACTAAGA CTCGTGAAGG TTCATTATAC	600
	TATCACGTAA CTGCTACTAA CGGTAGTGGT ATTAGTGGTT GGATTACGC TGGTAAGGGC	660
	TTCACTACTA CTGCTACTGG TACACAAGTA CTTGGTGGTC TGTCACCTGA TAAGTCAGTT	720
25	ACAGCAACCA ACGATAACAG TGTTAAGATT GTTTACCGTA CGACTGATGG CACTCAAGTT	780
	GGTTCTAACA CTTGGGTAAC TTCAACTGAT GGTACAAAGG CAGGTTCTAA GGTAAAGCAT	840
30	AAGGCCGCCG ATCAAACGTC TCTGAAGCC TACATCAATG CTAACAAGCC TAGCGGTTAC	900
	ACTGTAACCT ACCCTAATGC TGCAGATGCT ACCTATGGTA ACACAGTTTA CGCTACTGTT	960
	TCCCAAGCAG CTACTTCTAA GGTGCTTTG AAGGTCTCAG GGACTCCTGT TACTACTGCA	1020
35	TTGACTACAG CTGATGCTAA TGATAAGGTT GCAGCTAACG ATACCACTGC TAATGGTAGT	1080
	TCTGTGTCAG GCTCAACAGT CTATGCTGCT GGTACTAAGT TGGCTCAATT AACAACTGAC	1140
40	TTGACTGGTG AAAAGGGTCA AGTTGTGACA TTAAGTGCCA TCGATACTGA TTTGGGAAGC	1200
	GCTACGTTCA CTGGAACCTAC GACTTACTAT TCAGATCTTG GTAAAGCATA CCACTACACT	1260
	TACACTTACA ATAAGGACAG TGCTGCTTCT TCAATGCAA GTACCAATT TGGTTCAAAC	1320
45	GTCACTGGTA CTTTAACTGC TACCCTTGTT ATGGGTAAGT CTAAGTCTAC TGCTAACGGT	1380
	ACTACTGGT TCAAC	1395

(2) INFORMATION FOR SEQ ID NO: 7:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 105 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

55

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

5 Val Lys Thr Thr Asn Arg Gly Ser Val Tyr Tyr Arg Val Val Thr Met
 1 5 10 15
 Asp Gly Lys Tyr Arg Gly Tyr Val Tyr Gly Gly Lys Ser Asp Thr Ala
 20 25 30
 10 Phe Ala Gly Gly Ile Lys Ser Ala Glu Thr Thr Thr Lys Ala Asp Met
 35 40 45
 Pro Ala Arg Thr Thr Gly Phe Tyr Leu Thr Asp Thr Ser Lys Asn Thr
 50 55 60
 15 Leu Trp Thr Ala Pro Lys Tyr Thr Gln Tyr Lys Ala Ser Lys Val Ser
 65 70 75 80
 Leu Tyr Gly Val Ala Lys Asp Thr Lys Phe Thr Val Asp Gln Ala Ala
 85 90 95
 20 Thr Lys Thr Arg Glu Gly Ser Leu Tyr
 100 105

(2) INFORMATION FOR SEQ ID NO: 8:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 150 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 30 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

35 Val Lys Thr Thr Asn Arg Gly Ser Val Tyr Tyr Arg Val Val Thr Met
 1 5 10 15
 40 Asp Gly Lys Tyr Arg Gly Tyr Val Tyr Gly Gly Lys Ser Asp Thr Ala
 20 25 30
 Phe Ala Gly Gly Ile Lys Ser Ala Glu Thr Thr Thr Lys Ala Asp Met
 35 40 45
 45 Pro Ala Arg Thr Thr Gly Phe Tyr Leu Thr Asp Thr Ser Lys Asn Thr
 50 55 60
 Leu Trp Thr Ala Pro Lys Tyr Thr Gln Tyr Lys Ala Ser Lys Val Ser
 65 70 75 80
 50 Leu Tyr Gly Val Ala Lys Asp Thr Lys Phe Thr Val Asp Gln Ala Ala
 85 90 95
 55 Thr Lys Thr Arg Glu Gly Ser Leu Tyr Tyr His Val Thr Ala Thr Asn
 100 105 110
 Gly Ser Gly Ile Ser Gly Trp Ile Tyr Ala Gly Lys Gly Phe Ser Thr

34

	195		200		205
5	Asn Pro Asn Ala Ala Asp Ala Thr Tyr Gly Asn Thr Val Tyr Ala Thr				
	210		215		220
	Val Ser Gln Ala Ala Thr Ser Lys Val Ala Leu Lys Val Ser Gly Thr				
	225		230		235 240
10	Pro Val Thr Thr Ala Leu Thr Thr Ala Asp Ala Asn Asp Lys Val Ala				
		245		250	255
	Ala Asn Asp Thr Thr Ala Asn Gly Ser Ser Val Ala Gly Ser Thr Val				
		260		265	270
15	Tyr Ala Ala				
	275				

(2) INFORMATION FOR SEQ ID NO: 10:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 215 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: both

25 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

30	Lys Ser Tyr Ala Thr Ala Gly Ala Tyr Ser Thr Leu Lys Thr Asp Ala
	1 5 10 15
	Ala Thr Arg Asn Val Glu Ala Thr Gly Thr Asn Ala Leu Tyr Thr Lys
	20 25 30
35	Pro Gly Thr Val Lys Gly Ala Lys Val Val Ala Ser Lys Ala Thr Met
	35 40 45
40	Ala Lys Leu Ala Ser Ser Lys Lys Ser Ala Asp Tyr Phe Arg Ala Tyr
	50 55 60
	Gly Val Lys Thr Thr Asn Arg Gly Ser Val Tyr Tyr Arg Val Val Thr
	65 70 75 80
45	Met Asp Gly Lys Tyr Arg Gly Tyr Val Tyr Gly Gly Lys Ser Asp Thr
	85 90 95
	Ala Phe Ala Gly Gly Ile Lys Ser Ala Glu Thr Thr Thr Lys Ala Asp
	100 105 110
50	Met Pro Ala Arg Thr Thr Gly Phe Tyr Leu Thr Asp Thr Ser Lys Asn
	115 120 125
	Thr Leu Trp Thr Ala Pro Lys Tyr Thr Gln Tyr Lys Ala Ser Lys Val
	130 135 140
55	Ser Leu Tyr Gly Val Ala Lys Asp Thr Lys Phe Thr Val Asp Gln Ala

35

145 150 155 160

Ala Thr Lys Thr Arg Glu Gly Ser Leu Tyr Tyr His Val Thr Ala Thr
165 170 175

5 Asn Gly Ser Gly Ile Ser Gly Trp Ile Tyr Ala Gly Lys Gly Phe Ser
180 185 190

10 Thr Thr Ala Thr Gly Thr Gln Val Leu Gly Gly Leu Ser Thr Asp Lys
195 200 205

Ser Val Thr Ala Thr Asn Asp
210 215

15 (2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 270 amino acids

20 (B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Lys Ser Tyr Ala Thr Ala Gly Ala Tyr Ser Thr Leu Lys Thr Asp Ala
1 5 10 15

30 Ala Thr Arg Asn Val Glu Ala Thr Gly Thr Asn Ala Leu Tyr Thr Lys
20 25 30

35 Pro Gly Thr Val Lys Gly Ala Lys Val Val Ala Ser Lys Ala Thr Met
35 40 45

Ala Lys Leu Ala Ser Ser Lys Lys Ser Ala Asp Tyr Phe Arg Ala Tyr
50 55 60

40 Gly Val Lys Thr Thr Asn Arg Gly Ser Val Tyr Tyr Arg Val Val Thr
65 70 75 80

Met Asp Gly Lys Tyr Arg Gly Tyr Val Tyr Gly Gly Lys Ser Asp Thr
85 90 95

45 Ala Phe Ala Gly Gly Ile Lys Ser Ala Glu Thr Thr Thr Lys Ala Asp
100 105 110

50 Met Pro Ala Arg Thr Thr Gly Phe Tyr Leu Thr Asp Thr Ser Lys Asn
115 120 125

Thr Leu Trp Thr Ala Pro Lys Tyr Thr Gln Tyr Lys Ala Ser Lys Val
130 135 140

55 Ser Leu Tyr Gly Val Ala Lys Asp Thr Lys Phe Thr Val Asp Gln Ala
145 150 155 160

36

Ala Thr Lys Thr Arg Glu Gly Ser Leu Tyr Tyr His Val Thr Ala Thr
165 170 175

5 Asn Gly Ser Gly Ile Ser Gly Trp Ile Tyr Ala Gly Lys Gly Phe Ser
180 185 190

Thr Thr Ala Thr Gly Thr Gln Val Leu Gly Gly Leu Ser Thr Asp Lys
195 200 205

10 Ser Val Thr Ala Thr Asn Asp Asn Ser Val Lys Ile Val Tyr Arg Thr
210 215 220

Thr Asp Gly Thr Gln Val Gly Ser Asn Thr Trp Val Thr Ser Thr Asp
225 230 235 240

15 Gly Thr Lys Ala Gly Ser Lys Val Ser Asp Lys Ala Ala Asp Gln Thr
245 250 255

Ala Leu Glu Ala Tyr Ile Asn Ala Asn Lys Pro Ser Gly Tyr
260 265 270

20

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
- 25 (A) LENGTH: 465 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: both
- 30 (ii) MOLECULE TYPE: protein
- (x) PUBLICATION INFORMATION:
- 35 (A) AUTHORS: Vidgren, G
Palva, I
Pakkanen, R
Lounatmaa, K
Palva, A
- 40 (B) TITLE: S-Layer Protein Gene of Lactobacillus brevis:
Cloning by Polymerase Chain Reaction and
Determination of the Nucleotide sequence
- (C) JOURNAL: J. Bacteriol.
(D) VOLUME: 174
(E) ISSUE: 22
(F) PAGES: 7419-7427
45 (G) DATE: 1992
(K) RELEVANT RESIDUES IN SEQ ID NO: 12: FROM 1 TO 465
- (x) PUBLICATION INFORMATION:
- 50 (H) DOCUMENT NUMBER: WO 94/00581 A1
(I) FILING DATE: 24-JUN-1993
(J) PUBLICATION DATE: 06-JAN-1994
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
- 55 Met Gln Ser Ser Leu Lys Lys Ser Leu Tyr Leu Gly Leu Ala Ala Leu
1 5 10 15

	Ser	Phe	Ala	Gly	Val	Ala	Ala	Val	Ser	Thr	Thr	Ala	Ser	Ala	Lys	Ser	
				20					25						30		
5	Tyr	Ala	Thr	Ala	Gly	Ala	Tyr	Ser	Thr	Leu	Lys	Thr	Asp	Ala	Ala	Thr	
			35					40					45				
	Arg	Asn	Val	Glu	Ala	Thr	Gly	Thr	Asn	Ala	Leu	Tyr	Thr	Lys	Pro	Gly	
		50					55					60					
10	Thr	Val	Lys	Gly	Ala	Lys	Val	Val	Ala	Ser	Lys	Ala	Thr	Met	Ala	Lys	
		65				70					75					80	
	Leu	Ala	Ser	Ser	Lys	Lys	Ser	Ala	Asp	Tyr	Phe	Arg	Ala	Tyr	Gly	Val	
					85					90					95		
15	Lys	Thr	Thr	Asn	Arg	Gly	Ser	Val	Tyr	Tyr	Arg	Val	Val	Thr	Met	Asp	
				100					105						110		
20	Gly	Lys	Tyr	Arg	Gly	Tyr	Val	Tyr	Gly	Gly	Lys	Ser	Asp	Thr	Ala	Phe	
			115					120					125				
	Ala	Gly	Gly	Ile	Lys	Ser	Ala	Glu	Thr	Thr	Thr	Lys	Ala	Asp	Met	Pro	
			130				135						140				
25	Ala	Arg	Thr	Thr	Gly	Phe	Tyr	Leu	Thr	Asp	Thr	Ser	Lys	Asn	Thr	Leu	
		145				150				155						160	
	Trp	Thr	Ala	Pro	Lys	Tyr	Thr	Gln	Tyr	Lys	Ala	Ser	Lys	Val	Ser	Leu	
				165						170					175		
30	Tyr	Gly	Val	Ala	Lys	Asp	Thr	Lys	Phe	Thr	Val	Asp	Gln	Ala	Ala	Thr	
				180					185					190			
35	Lys	Thr	Arg	Glu	Gly	Ser	Leu	Tyr	Tyr	His	Val	Thr	Ala	Thr	Asn	Gly	
			195					200					205				
	Ser	Gly	Ile	Ser	Gly	Trp	Ile	Tyr	Ala	Gly	Lys	Gly	Phe	Ser	Thr	Thr	
		210					215					220					
40	Ala	Thr	Gly	Thr	Gln	Val	Leu	Gly	Gly	Leu	Ser	Thr	Asp	Lys	Ser	Val	
					230					235					240		
	Thr	Ala	Thr	Asn	Asp	Asn	Ser	Val	Lys	Ile	Val	Tyr	Arg	Thr	Thr	Asp	
				245						250					255		
45	Gly	Thr	Gln	Val	Gly	Ser	Asn	Thr	Trp	Val	Thr	Ser	Thr	Asp	Gly	Thr	
			260						265					270			
50	Lys	Ala	Gly	Ser	Lys	Val	Ser	Asp	Lys	Ala	Ala	Asp	Gln	Thr	Ala	Leu	
			275					280					285				
	Glu	Ala	Tyr	Ile	Asn	Ala	Asn	Lys	Pro	Ser	Gly	Tyr	Thr	Val	Thr	Asn	
			290				295					300					
55	Pro	Asn	Ala	Ala	Asp	Ala	Thr	Tyr	Gly	Asn	Thr	Val	Tyr	Ala	Thr	Val	
		305				310					315					320	

38

Ser Gln Ala Ala Thr Ser Lys Val Ala Leu Lys Val Ser Gly Thr Pro
 325 330 335
 Val Thr Thr Ala Leu Thr Thr Ala Asp Ala Asn Asp Lys Val Ala Ala
 340 345 350
 Asn Asp Thr Thr Ala Asn Gly Ser Ser Val Ala Gly Ser Thr Val Tyr
 355 360 365
 Ala Ala Gly Thr Lys Leu Ala Gln Leu Thr Thr Asp Leu Thr Gly Glu
 370 375 380
 Lys Gly Gln Val Val Thr Leu Thr Ala Ile Asp Thr Asp Leu Glu Asp
 385 390 395 400
 Ala Thr Phe Thr Gly Thr Thr Thr Tyr Tyr Ser Asp Leu Gly Lys Ala
 405 410 415
 Tyr His Tyr Thr Tyr Thr Tyr Asn Lys Asp Ser Ala Ala Ser Ser Asn
 420 425 430
 Ala Ser Thr Gln Phe Gly Ser Asn Val Thr Gly Thr Leu Thr Ala Thr
 435 440 445
 Leu Val Met Gly Lys Ser Thr Ala Thr Ala Asn Gly Thr Thr Trp Phe
 450 455 460
 Asn
 465

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 209 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Ser Val Thr Ala Thr Asn Asp Asn Ser Val Lys Ile Val Tyr Arg Thr
 1 5 10 15
 Thr Asp Gly Thr Gln Val Gly Ser Asn Thr Trp Val Thr Ser Thr Asp
 20 25 30
 Gly Thr Lys Ala Gly Ser Lys Val Ser Asp Lys Ala Ala Asp Gln Thr
 35 40 45
 Ala Leu Glu Ala Tyr Ile Asn Ala Asn Lys Pro Ser Gly Tyr Thr Val
 50 55 60
 Thr Asn Pro Asn Ala Ala Asp Ala Thr Tyr Gly Asn Thr Val Tyr Ala
 65 70 75 80

39

	Thr	Val	Ser	Gln	Ala	Ala	Thr	Ser	Lys	Val	Ala	Leu	Lys	Val	Ser	Gly	
				85						90						95	
5	Thr	Pro	Val	Thr	Thr	Ala	Leu	Thr	Thr	Ala	Asp	Ala	Asn	Asp	Lys	Val	
				100					105						110		
	Ala	Ala	Asn	Asp	Thr	Thr	Ala	Asn	Gly	Ser	Ser	Val	Ala	Gly	Ser	Thr	
			115					120					125				
10	Val	Tyr	Ala	Ala	Gly	Thr	Lys	Leu	Ala	Gln	Leu	Thr	Thr	Asp	Leu	Thr	
		130					135					140					
	Gly	Glu	Lys	Gly	Gln	Val	Val	Thr	Leu	Thr	Ala	Ile	Asp	Thr	Asp	Leu	
15		145			150						155					160	
	Glu	Asp	Ala	Thr	Phe	Thr	Gly	Thr	Thr	Thr	Tyr	Tyr	Ser	Asp	Leu	Gly	
				165						170					175		
	Lys	Ala	Tyr	His	Tyr	Thr	Tyr	Thr	Tyr	Asn	Lys	Asp	Ser	Ala	Ala	Ser	
20				180					185						190		
	Ser	Asn	Ala	Ser	Thr	Gln	Phe	Gly	Ser	Asn	Val	Thr	Gly	Thr	Leu	Thr	
			195					200					205				
25	Ala																
(2) INFORMATION FOR SEQ ID NO: 14:																	
30	(i) SEQUENCE CHARACTERISTICS:																
	(A) LENGTH: 627 base pairs																
	(B) TYPE: nucleic acid																
	(C) STRANDEDNESS: both																
	(D) TOPOLOGY: both																
35	(ii) MOLECULE TYPE: DNA (genomic)																
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:																
40	TCAGTTACAG	CAACCAACGA	TAACAGTGTT	AAGATTGTTT	ACCGTACGAC	TGATGGCACT											60
	CAAGTTGGTT	CTAACACTTG	GGTAACTTCA	ACTGATG GTA	CAAAGGCAGG	TTCTAAGSTA											120
45	AGCGATAAGG	CCGCCGATCA	AAC TGCTCTT	GAAGCCTACA	TCAATGCTAA	CAAGCCTAGC											180
	GGTTACACTG	TAACTAACC	TAATGCTGCA	GATGCTACCT	ATG GTAACAC	AGTTACGCT											240
	ACTGTTTCCC	AAGCAGCTAC	TTCTAAGGTC	GCTTTGAAGG	TCTCAGGGAC	TCCTGTTACT											300
50	ACTGCATTGA	CTACAGCTGA	TGCTAATGAT	AAGGTTGCAG	CTAACGATAC	CACTGCTAAT											360
	GGTAGTTC TG	TTGCAGGCTC	AACAGTCTAT	GCTGCTGGTA	CTAAGTTGGC	TCAATTAACA											420
55	ACTGACTTGA	CTGGTGAAAA	GGGTCAAGTT	GTCACATTAA	CTGCCATCGA	TACTGATT TG											480
	GAAGACGCTA	CGTTCACTGG	AAC TACGACT	TACTATT CAG	ATCTTG GTA	AGCATACCAC											540

TACACTTACA CTTACAATAA GGACAGTGCT GCTTCTTCAA ATGCAAGTAC CCAATTGGT 600
TCAACGTC A CTGGTACTTT AACTGCT 627

5 (2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1680 base pairs
(B) TYPE: nucleic acid
10 (C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: DNA (genomic)

15

(ix) FEATURE:

- (A) NAME/KEY: CDS
(B) LOCATION:189..1583

20

(ix) FEATURE:

- (A) NAME/KEY: misc_signal
(B) LOCATION:189..278

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

25

TCCAACGACA ATCAGAGCGT AATCCTTGTA TCTCCTTAAG GAAATCGCTA TACTTATCTT 60

CGTAGTTAGG GATAGCTGA TCGGGTCCGC TAATGTTATG AAATAAAATT CTTAACAAAA 120

30

CGCGTAACTT CGGTTACTT ATTCTTGCTT GATAAATTAC ATATTTTATG TTTGAGGAA 180

GAAAGATT ATG CAA TCA AGT TTA AAG AAA TCT CTT TAC TTG GGC CTT GCC 230
Met Gln Ser Ser Leu Lys Lys Ser Leu Tyr Leu Gly Leu Ala
1 5 10

35

GCA TTG AGC TTT GCT GGT GTT GCT GCC GTT TCA ACG ACT GCT TCA GCT 278
Ala Leu Ser Phe Ala Thr Val Ala Ala Val Ser Thr Thr Ala Ser Ala
15 20 25 30

40

AAG TCA TAC GCT ACT GCA GGT GCC TAT TCA ACG TTA AAG ACG GAC GCT 326
Lys Ser Tyr Ala Thr Ala Gly Ala Tyr Ser Thr Leu Lys Thr Asp Ala
35 40 45

45

GCT ACT CGT AAC GTC GAA GCT ACT GGT ACT AAC GCT TTA TAC ACG AAG 374
Ala Thr Arg Asn Val Glu Ala Thr Gly Thr Asn Ala Leu Tyr Thr Lys
50 55 60

50

CCA GGT ACT GTT AAG GGT GCT AAG GTT GTC GCT TCT TCT AAG GCT ACT ATG 422
Pro Gly Thr Val Lys Gly Ala Lys Val Val Ala Ser Lys Ala Thr Met
65 70 75

55

GCT AAG TTA GCT TCT TCA AAG AAG TCA GCT GAC TAC TTC CGT GCT TAC 470
Ala Lys Leu Ala Ser Ser Lys Lys Ser Ala Asp Tyr Phe Arg Ala Tyr
80 85 90

GGT GTT AAG ACC ACT AAC CGT GGT TCA GTT TAC TAC CGT GTT GTA ACG 518
Gly Val Lys Thr Thr Asn Arg Gly Ser Val Tyr Tyr Arg Val Val Thr

41

	95	100	105	110	
	ATG GAT GGC AAG TAC CGT GGT TAC GTT TAT GGT GGC AAG TCT GAC ACT				566
5	Met Asp Gly Lys Tyr Arg Gly Tyr Val Tyr Gly Gly Lys Ser Asp Thr	115	120	125	
	GCC TTT GCT GGT GGT ATC AAG TCT GCT GAA ACG ACT ACT AAG GCT GAT				614
	Ala Phe Ala Gly Lys Ile Lys Ser Ala Glu Thr Thr Thr Lys Ala Asp	130	135	140	
10	ATG CCT GCA CGT ACT ACT GGG TTC TAC TTA ACT GAC ACT TCA AAG AAC				662
	Met Pro Ala Arg Thr Thr Gly Phe Tyr Leu Thr Asp Thr Ser Lys Asn	145	150	155	
15	ACT CTT TGG ACG GCT CCT AAG TAC ACT CAA TAC AAG GCA AGT AAA GTT				710
	Thr Leu Trp Thr Ala Pro Lys Tyr Thr Gln Tyr Lys Ala Ser Lys Val	160	165	170	
20	AGC CTT TAT GGT GTT GCT AAG GAC ACC AAG TTT ACT GTA GAT CAG GCT				758
	Ser Leu Tyr Gly Val Ala Lys Asp Thr Lys Phe Thr Val Asp Gln Ala	175	180	185	190
25	GCT ACT AAG ACT CGT GAA GGT TCA TTA TAC TAT CAC GTA ACT GCT ACT				806
	Ala Thr Lys Thr Arg Glu Gly Ser Leu Tyr Tyr His Val Thr Ala Thr	195	200	205	
30	AAC GGT AGT GGT ATT AGT GGT TGG ATT TAC GCT GGT AAG GGC TTC AGT				854
	Asn Gly Ser Gly Ile Ser Gly Trp Ile Tyr Ala Gly Lys Gly Phe Ser	210	215	220	
	ACT ACT GCT ACT GGT ACA CAA GTA CIT GGT GGT CTG TCA ACT GAT AAG				902
	Thr Thr Ala Thr Gly Thr Gln Val Leu Gly Gly Leu Ser Thr Asp Lys	225	230	235	
35	TCA GTT ACA GCA ACC AAC GAT AAC AGT GTT AAG ATT GTT TAC CGT ACG				950
	Ser Val Thr Ala Thr Asn Asp Asn Ser Val Lys Ile Val Tyr Arg Thr	240	245	250	
40	ACT GAT GGC ACT CAA GTT GGT TCT AAC ACT TGG GTA ACT TCA ACT GAT				998
	Thr Asp Gly Thr Gln Val Gly Ser Asn Thr Trp Val Thr Ser Thr Asp	255	260	265	270
45	GGT ACA AAG GCA GGT TCT AAG GTA AGC GAT AAG GCC GCC GAT CAA ACT				1046
	Gly Thr Lys Ala Gly Ser Lys Val Ser Asp Lys Ala Ala Asp Gln Thr	275	280	285	
50	GCT CTT GAA GCC TAC ATC AAT GCT AAC AAG CCT AGC GGT TAC ACT GTA				1094
	Ala Leu Glu Ala Tyr Ile Asn Ala Asn Lys Pro Ser Gly Tyr Thr Val	290	295	300	
	ACT AAC CCT AAT GCT GCA GAT GCT ACC TAT GGT AAC ACA GTT TAC GCT				1142
	Thr Asn Pro Asn Ala Ala Asp Ala Thr Tyr Gly Asn Thr Val Tyr Ala	305	310	315	
55	ACT GTT TCC CAA GCA GCT ACT TCT AAG GTC GCT TTG AAG GTC TCA GGG				1190
	Thr Val Ser Gln Ala Ala Thr Ser Lys Val Ala Leu Lys Val Ser Gly	320	325	330	

42

	ACT CCT GTT ACT ACT GCA TTG ACT ACA GCT GAT GCT AAT GAT AAG GTT	1238
	Thr Pro Val Thr Thr Ala Leu Thr Thr Ala Asp Ala Asn Asp Lys Val	
	335 340 345 350	
5	GCA GCT AAC GAT ACC ACT GCT AAT GGT AGT TCT GTT GCA GGC TCA ACA	1286
	Ala Ala Asn Asp Thr Thr Ala Asn Gly Ser Ser Val Ala Gly Ser Thr	
	355 360 365	
	GTC TAT GCT GCT GGT ACT AAG TTG GCT CAA TTA ACA ACT GAC TTG ACT	1334
10	Val Tyr Ala Ala Gly Thr Lys Leu Ala Gln Leu Thr Thr Asp Leu Thr	
	370 375 380	
	GGT GAA AAG GGT CAA GTT GTC ACA TTA ACT GCC ATC GAT ACT GAT TTG	1382
15	Gly Glu Lys Gly Gln Val Val Thr Leu Thr Ala Ile Asp Thr Asp Leu	
	385 390 395	
	GAA GAC GCT ACG TTC ACT GGA ACT ACG ACT TAC TAT TCA GAT CTT GGT	1430
20	Glu Asp Ala Thr Phe Thr Gly Thr Thr Tyr Tyr Ser Asp Leu Gly	
	400 405 410	
	AAA GCA TAC CAC TAC ACT TAC ACT TAC AAT AAG GAC AGT GCT TCT	1478
	Lys Ala Tyr His Tyr Thr Tyr Thr Tyr Asn Lys Asp Ser Ala Ala Ser	
	415 420 425 430	
25	TCA AAT GCA AGT ACC CAA TTT GGT TCA AAC GTC ACT GGT ACT TTA ACT	1526
	Ser Asn Ala Ser Thr Gln Phe Gly Ser Asn Val Thr Gly Thr Leu Thr	
	435 440 445	
	GCT ACC CTT GTT ATG GGT AAG TCT ACT GCT ACT GCT AAC GGT ACT ACT	1574
30	Ala Thr Leu Val Met Gly Lys Ser Thr Ala Thr Ala Asn Gly Thr Thr	
	450 455 460	
	TGG TTC AAC TAATAATTAT TATTTAGGTG AGCTTTGTTG ATAAAAAGGT	1623
35	Trp Phe Asn	
	465	
	CTTTTCAACG TTTATGTTGG GGAGACCTTT TTATATTGAA AAAATTAGGC CTTTGT	1680
40	(2) INFORMATION FOR SEQ ID NO: 16:	
	(i) SEQUENCE CHARACTERISTICS:	
	(A) LENGTH: 465 amino acids	
	(B) TYPE: amino acid	
45	(D) TOPOLOGY: linear	
	(ii) MOLECULE TYPE: protein	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:	
50	Met Gln Ser Ser Leu Lys Lys Ser Leu Tyr Leu Gly Leu Ala Ala Leu	
	1 5 10 15	
	Ser Phe Ala Gly Val Ala Ala Val Ser Thr Thr Ala Ser Ala Lys Ser	
55	20 25 30	
	Tyr Ala Thr Ala Gly Ala Tyr Ser Thr Leu Lys Thr Asp Ala Ala Thr	

43

	35		40		45
	Arg Asn Val Glu Ala Thr Gly Thr Asn Ala Leu Tyr Thr Lys Pro Gly				
	50		55		60
5	Thr Val Lys Gly Ala Lys Val Val Ala Ser Lys Ala Thr Met Ala Lys				
	65		70		75
	Leu Ala Ser Ser Lys Lys Ser Ala Asp Tyr Phe Arg Ala Tyr Gly Val				
10			85		90
	Lys Thr Thr Asn Arg Gly Ser Val Tyr Tyr Arg Val Val Thr Met Asp				
			100		105
					110
15	Gly Lys Tyr Arg Gly Tyr Val Tyr Gly Gly Lys Ser Asp Thr Ala Phe				
			115		120
	Ala Gly Gly Ile Lys Ser Ala Glu Thr Thr Thr Lys Ala Asp Met Pro				
			130		135
					140
20	Ala Arg Thr Thr Gly Phe Tyr Leu Thr Asp Thr Ser Lys Asn Thr Leu				
			145		150
					155
	Trp Thr Ala Pro Lys Tyr Thr Gln Tyr Lys Ala Ser Lys Val Ser Leu				
25			165		170
					175
	Tyr Gly Val Ala Lys Asp Thr Lys Phe Thr Val Asp Gln Ala Ala Thr				
			180		185
					190
30	Lys Thr Arg Glu Gly Ser Leu Tyr Tyr His Val Thr Ala Thr Asn Gly				
			195		200
	Ser Gly Ile Ser Gly Trp Ile Tyr Ala Gly Lys Gly Phe Ser Thr Thr				
			210		215
					220
35	Ala Thr Gly Thr Gln Val Leu Gly Gly Leu Ser Thr Asp Lys Ser Val				
			225		230
					235
	Thr Ala Thr Asn Asp Asn Ser Val Lys Ile Val Tyr Arg Thr Thr Asp				
40			245		250
					255
	Gly Thr Gln Val Gly Ser Asn Thr Trp Val Thr Ser Thr Asp Gly Thr				
			260		265
					270
45	Lys Ala Gly Ser Lys Val Ser Asp Lys Ala Ala Asp Gln Thr Ala Leu				
			275		280
	Glu Ala Tyr Ile Asn Ala Asn Lys Pro Ser Gly Tyr Thr Val Thr Asn				
			290		295
					300
50	Pro Asn Ala Ala Asp Ala Thr Tyr Gly Asn Thr Val Tyr Ala Thr Val				
			305		310
					315
	Ser Gln Ala Ala Thr Ser Lys Val Ala Leu Lys Val Ser Gly Thr Pro				
			325		330
					335
55	Val Thr Thr Ala Leu Thr Thr Ala Asp Ala Asn Asp Lys Val Ala Ala				

44

		340				345						350
		Asn Asp Thr Thr Ala Asn Gly Ser Ser Val Ala Gly Ser Thr Val Tyr										
		355				360						365
5		Ala Ala Gly Thr Lys Leu Ala Gln Leu Thr Thr Asp Leu Thr Gly Glu										
		370				375						380
10		Lys Gly Gln Val Val Thr Leu Thr Ala Ile Asp Thr Asp Leu Glu Asp										
		385				390						400
		Ala Thr Phe Thr Gly Thr Thr Thr Tyr Tyr Ser Asp Leu Gly Lys Ala										
						405						415
15		Tyr His Tyr Thr Tyr Thr Tyr Asn Lys Asp Ser Ala Ala Ser Ser Asn										
						420						430
		Ala Ser Thr Gln Phe Gly Ser Asn Val Thr Gly Thr Leu Thr Ala Thr										
						435						445
20		Leu Val Met Gly Lys Ser Thr Ala Thr Ala Asn Gly Thr Thr Trp Phe										
						450						460
		Asn										
25		465										

What we claim is:

1. A DNA molecule encoding a polypeptide capable of binding to human and/or animal
epithelial cell types, said DNA molecule being selected from the group consisting of:
 - (a) DNA molecules having at least the partial coding sequences of any one of SEQ ID NO. 1, SEQ ID NO. 2, SEQ ID NO. 3, SEQ ID NO. 4, SEQ ID NO 5 and/or SEQ ID NO. 6 excluding the full length SEQ ID NO 6;
 - (b) DNA molecules encoding a polypeptide having at least the partial amino acid sequences of any one of SEQ ID NO. 7, SEQ ID NO. 8, SEQ ID NO.9, SEQ ID NO. 10, SEQ ID NO. 11 and/or SEQ ID NO 12, excluding the full length SEQ ID NO 12;
 - (c) DNA molecules the coding sequences of which differ from the coding sequence of a nucleic acid molecule of (a) or (b) due to the degeneracy of the genetic code;
 - (d) DNA molecules hybridizing under stringent conditions to a molecule of (a), (b) and/or (c) excluding the full-length SEQ ID NO 6 ; and
 - (e) DNA molecules encoding a polypeptide capable of binding to human and/or animal epithelial cell types and having an amino acid sequence which shows at least 40 % identity to a sequence contained in (b) excluding the full length SEQ ID NO 12.
2. The DNA molecule of claim 1 encoding a polypeptide capable of binding to human and/or animal epithelial cell types, wherein said polypeptide is capable of binding to intestinal, urogenital and/or endothelial cells.
3. The DNA molecule of claim 1 or 2, which originates from the DNA molecule encoding *Lactobacillus brevis* S-layer SlpA protein.
4. A vector containing a DNA molecule of any one of claim 1, 2 or 3.
5. The vector of claim 4, in which the DNA molecule is operably linked to expression and optionally to secretion control sequences allowing expression in prokaryotic or eukaryotic

host cells.

6. A host cell transformed with a DNA molecule of any one of claims 1, 2 or 3 or with a vector of claim 4 or 5.

5

7. A method of constructing a host cell capable of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cell types, comprising transforming the cell with at least one DNA molecule selected from the group consisting of:

- 10 (a) DNA molecules having at least the partial coding sequences of any one of SEQ ID NO. 1, SEQ ID NO. 2, SEQ ID NO. 3, SEQ ID NO. 4, SEQ ID NO. 5, and/or SEQ ID NO. 6;
- (b) DNA molecules encoding a polypeptide having at least the partial amino acid sequences of any one of SEQ ID NO. 7, SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 10, SEQ ID NO. 11 and/or SEQ ID NO. 12 ;
- 15 (c) DNA molecules, the coding sequence of which differ from the coding sequence of a DNA molecule of (a) or (b) due to the degeneracy of the genetic code;
- (d) DNA molecules hybridizing under stringent conditions to a molecule of (a), (b) and/or (c); and
- (e) DNA molecules encoding a polypeptide capable of binding to human and/or animal
- 20 epithelial cell types, like intestinal, urogenital and/or endothelial cell types and having an amino acid sequence which shows at least 40 % identity to a sequence contained in (b).

8. A host cell constructed by the method of claim 7.

- 25 9. The host cell of any one of claims 6 or 8, which has probiotic effects.

10. The host cell of claim 9, wherein the probiotic effects have been enhanced by genetic means.

- 30 11. The host cell of claim 9 or 10, which belongs to lactic acid bacteria or bifidobacteria.

12. The host cell of any one of claims 6 or 8 to 11, wherein the host cell is a vaccine carrier.

13. The host cell of any one of claims 6 or 8 to 12, wherein the host cell has been genetically modified to carry at least one of the factors selected from the group consisting of an antigen, an epitope, a single chain antibody, an enzyme, a pharmaceutical protein and a toxin.

5

14. A method of constructing a polypeptide capable of binding to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cell types, comprising modifying the gene encoding the polypeptide with a DNA molecule of claim 1 or 7.

10 15. A gene encoding a preselected protein, wherein the gene encoding the protein is genetically modified to bind to human and/or animal epithelial cell types, like intestinal, urogenital and/or endothelial cell types with at least one DNA molecule selected from the group consisting of any of the sequences of claim 1 or 7.

15 16. The gene of claim 15, wherein the preselected protein is selected from the group consisting of an antigen, an epitope, a single chain antibody, an enzyme, a pharmaceutical protein and a toxin.

17. A polypeptide encoded by a DNA molecule of claim 1 or 15.

20

18. A *Lactobacillus brevis* strain for carrying preselected factors/ properties to human and/or animal epithelial cells or cell surfaces, wherein the strain has the capability of binding to human and/or animal epithelial cell types and wherein the gene encoding the S-layer SlpA protein is genetically modified to carry at least one of the factors/properties
25 selected from the group consisting of an antigen, an epitope, a single chain antibody, an enzyme, a pharmaceutical protein and a toxin.

19. A *Lactobacillus brevis* strain for carrying preselected factors/ properties to human and/or animal epithelial cells or cell surfaces, wherein the strain has the capability of
30 binding to human and/or animal epithelial cell types and wherein the strain is genetically modified to have improved probiotic effects.

20. A *Lactobacillus brevis* S-layer SlpA protein for carrying preselected factors/properties to human and/or animal epithelial cells or cell surfaces, wherein the protein has the capability of binding to human and/or animal epithelial cell types and wherein the gene encoding the S-layer SlpA protein is genetically modified to carry at least one of the
- 5 factors/properties selected from the group consisting of an antigen, an epitope, a single chain antibody, an enzyme, a pharmaceutical protein and a toxin.
21. A method for carrying preselected factors/ properties to human and/or animal epithelial cells or cell surfaces by using a *Lactobacillus brevis* strain, wherein the strain has the
- 10 capability of binding to human and/or animal epithelial cell types and wherein the gene encoding the S-layer SlpA protein is genetically modified to carry at least one of the factors/properties selected from the group consisting of an antigen, an epitope, a single chain antibody, an enzyme, a pharmaceutical protein and a toxin.
- 15 22. A method for carrying preselected factors/ properties to human and/or animal epithelial cells or cell surfaces by using a *Lactobacillus brevis* strain, wherein the strain has the capability of binding to human and/or animal epithelial cell types and wherein the strain is genetically modified to have improved probiotic effects.
- 20 23. A method for carrying preselected factors/ properties to human and/or animal epithelial cells or cell surfaces by using the *Lactobacillus brevis* S-layer SlpA protein, wherein the protein has the capability of binding to human and/or animal epithelial cell types and wherein the gene encoding the S-layer SlpA protein is genetically modified to carry at least one of the factors/properties selected from the group consisting of an antigen, an
- 25 epitope, a single chain antibody, an enzyme, a pharmaceutical protein and a toxin.
24. The use of the host cells of any one of claims 6 or 8 to 13 for excluding pathogens in the cell surfaces of the gastrointestinal or urogenital tract of humans and/ or animals.

Fig. 1A



Fig. 1B



2/8




Fig. 2B




Fig. 2A

3/8

Expression of *slpA* fragments as fusions to *fliC*




Fig. 3A

**Binding to
Int 407 cells**

31	████████████████████	300	+
31	████████████████	245	+
96	████████████████████████████	370	+
96	██████████████	245	+
96	██████████	200	+
239	████████████████████████████████	447	-

Fig. 3B



Fig. 4A




Fig. 4B



Fig. 4C



Fig. 4D



Fig. 4E



Fig. 4F



Fig. 4G



Fig. 4H

5/8

| | | | |
 TCCAACGACAATCAGAGCGTAATCCTTGTATCTCCTTAAGGAAATCGCTATACTTATCTT

| | | | |
 CGTAGTTAGGGGATAGCTGATCGGGTCCGCTAATGTTATGAAATAAAATCTTAAACAAA

| | | | |
 GCGCTAACTTCGGTTATACTATTCTTGCTTGATAAAATTACATATTTTATGTTTGGAGGAA

1
 ↓
 GAAAGATTATGCAATCAAGTTTAAAGAAATCTCTTTACTTGGGCCTTGCCGCATTGAGCT
 (M) Q S S L K K S L Y L G L A A L S
 (slpA coding region
 1

| | | | |
 91
 ↓
 TTGCTGGTGTGCTGCCGTTTCAACGACTGCTTCAGCTAAGTCATACGCTACTGCAGGTG
 F A G V A A V S T T A S A (K) S Y A T A G
 31

| | | | |
 CCTATTCAACGTAAAGACGGACGCTGCTACTCGTAACGTCGAAGCTACTGGTACTAAGC
 A Y S T L K T D A A T R N V E A T G T N

| | | | |
 CTTTATACACGAAGCCAGGTACTGTTAAGGGTGCTAAGGTTGTCGCTTCTAAGGCTACTA
 A L Y T K P G T V K G A K V V A S K A T

| | | | | |
 286
 ↓
 TGGCTAAGTTAGCTTCTTCAAAGAAGTCAGCTGACTACTTCCGTGCTTACGGTGTAAAGA
 M A K L A S S K K S A D Y F R A Y G (V) K
 96

Fig. 5A

6/8

CCACTAACCGTGGTTCAGTTTACTACCGTGTGTAAACGATGGCAAGTACCGTGGTT
T T N R G S V Y Y R V V T M D G K Y R G

ACGTTTATGGTGGCAAGTCTGACACTGCCTTTGCTGGTGGTATCAAGTCTGCTGAAACGA
Y V Y G G K S D T A F A G G I K S A E T

CTACTAAGGCTGATATGCTGCACGTACTACTGGGTTCTACTTAAGTACACTTCAAAGA
T T K A D M P A R T T G F Y L T D T S K

ACACTCTTTGGACGGCTCCTAAGTACACTCAATACAAGGCAAGTAAAGTTAGCCTTTATG
N T L W T A P K Y T Q Y K A S K V S L Y

GTGTGCTAAGGACACCAAGTTTACTGTAGATCAGGCTGCTACTAAGACTCGTGAAGGT
G V A K D T K F T V D Q A A T K T R E G

600
↓
CATTATACTATCACGTAAGTCTACTAAGGTTAGTGGTATTAGTGGTGGATTACGCTG
S L (Y) Y H V T A T N G S G I S G W I Y A
200

GTAAGGGCTTCAGTACTACTGCTACTGGTACACAAGTACTGGTGGTCTGTCAACTGATA
G K G F S T T A T G T Q V L G G L S T D

715 735
↓ ↓
AGTCAGTTACAGCAACCAACGATAACAGTGTTAAGATTGTTTACCGTACGACTGATGGCA
K (S) V T A T N (D) N S V K I V Y R T T D G
239 245

Fig. 5B

7/8

CTCAAGTTGGTTCTAACACTTGGGTAACCTCAACTGATGGTACAAAGGCAGGTTCTAAGG
T Q V G S N T W V T S T D G T K A G S K

TAAAGCGATAAGGCCGCCGATCAAACCTGCTCTTGAAGCCTACATCAATGCTAACAAGCCTA
V S D K A A D Q T A L E A Y I N A N K P

900 ↓
GCGGTTACACTGTAACCTAACCCCTAATGCTGCAGATGCTACCTATGGTAACACAGTTTACG
S G (Y) T V T N P N A A D A T Y G N T V Y
300

CTACTGTTTCCCAAGCAGCTACTTCTAAGGTCGCTTTGAAGGTCTCAGGGACTCCTGTTA
A T V S Q A A T S K V A L K V S G T P V

CTACTGCATTGACTACAGCTGATGCTAATGATAAGGTTGCAGCTAACGATACCAGCTGCTA
T T A L T T A D A N D K V A A N D T T A

1110 ↓
ATGGTAGTTCGTTCAGGCTCAACAGTCTATGCTGCTGGTACTAAGTTGGCTCAATTA
N G S S V A G S T V Y A (A) G T K L A Q L
370

CAACTGACTTGACTGGTGAAAAGGGTCAAGTTGTACATTAACTGCCATCGATACTGATT
T T D L T G E K G Q V V T L T A I D T D

TGGAAGACGCTACGTTCACTGGAACCTACGACTTACTATTGAGATCTTGGTAAGCATACC
L E D A T F T G T T T Y Y S D L G K A Y

Fig. 5C

8/8

| | | | |
 ACTACACTTACACTTACAATAAGGACAGTGTCTTCTTCAAATGCAAGTACCCAATTTG
 H Y T Y T Y N K D S A A S S N A S T Q F

| | 1341 | | |
 ↓
 GTTCAAACGTCACTGGTACTTTAACTGCTACCCCTTGTATGGGTAAGTCTACTGCTACTG
 G S N V T G T L T (A) T L V M G K S T A T
 447

| | 1395 | | |
 ↓
 CTAACGGTACTACTTGGTTCAACTAATAATTATTATTAGGTGAGCTTTGTTGATAAAAA
 A N G T T W F (N)
 465

| | | | |
 GGCTTTTCAACGTTTATGTTGGGGAGACCTTTTATATTGAAAAAATTAGGCCTTTGTT

Fig. 5D

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 99/00290

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07K 14/325

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, PATENT ABSTRACTS OF JAPAN, STRAND, CASEARCH, BIOSIS, MEDLINE, SCISEARCH
H, LIFE SCIENCES COLLECTION

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FEMS Microbiology Reviews, Volume 20, 1997, Hubert Bahl et al, "IV. Molecular biology of S-layers", page 47 - page 98, see pages 82-83	18-23
A	--	1-17,24
A	WO 9400581 A1 (VIAGEN OY), 6 January 1994 (06.01.94)	1-24
A	Journal of Bacteriology, Volume 174, No 22, 1992, Gabriele vidgrén et al, "S-Layer Protein Gene of Lactobacillus brevis: Cloning by Polymerase ChainReaction and Determinaiton of the Nucleotide Sequence" page 7419 - page 7427	1-24
	--	

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

26 July 1999

28-07-1999

Name and mailing address of the ISA/

Authorized officer

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Hampus Rystedt/Eö

Facsimile No. +46 8 666 02 86

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 99/00290

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Journal of Applied Bacteriology, Volume 74, 1993, C. Schneitz et al, "Adhesion of Lactobacillus acidophilus to avian intestinal epithelial cells mediated by the crystalline bacterial cell surface layer (S-layer)" page 290 - page 294 --	1-24
A	Chemical Abstracts, Volume 126, No 13, 31 March 1997 (31.03.97), (Columbus, Ohio, USA), Savijoki, Kirsi et al, "High level heterologous protein production in Lactococcus and Lactobacillus using a new secretion system based on the Lactobacillus brevis S-layer signals", page 1, THE ABSTRACT No 167032, Gene 1997, 186 (2), 255-262 -- -----	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

01/07/99

PCT/FI 99/00290

Patent document
cited in search reportPublication
datePatent family
member(s)Publication
date

WO 9400581 A1

06/01/94

AU 4329793 A

24/01/94

INTERNATIONAL SEARCH REPORT

International application No.
PCT/FI 99/00290

A. CLASSIFICATION OF SUBJECT MATTER		
IPC6: C07K 14/325 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
WPI, PATENT ABSTRACTS OF JAPAN, STRAND, CASESEARCH, BIOSIS, MEDLINE, SCISEARCH H, LIFE SCIENCES COLLECTION		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FEMS Microbiology Reviews, Volume 20, 1997, Hubert Bahl et al, "IV. Molecular biology of S-layers", page 47 - page 98, see pages 82-83	18-23
A	--	1-17,24
A	WO 9400581 A1 (VIAGEN OY), 6 January 1994 (06.01.94)	1-24
A	Journal of Bacteriology, Volume 174, No 22, 1992, Gabriele vidgrén et al, "S-Layer Protein Gene of Lactobacillus brevis: Cloning by Polymerase ChainReaction and Determination of the Nucleotide Sequence" page 7419 - page 7427	1-24
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
26 July 1999		28-07-1999
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer Hampus Rystedt/E6 Telephone No. +46 8 782 25 00